Argyropoulos, Paul[Argyropoulos.Paul@epa.gov]; Lie, Sharyn[Lie.Sharyn@epa.gov]; To:

Korotney, David[korotney.david@epa.gov]

David DeRamus[david.deramus@bateswhite.com];

marc.chupka@brattle.com[marc.chupka@brattle.com]; Lindsay Fitzgerald[lfitzgerald@biodiesel.org];

Anne Steckel[asteckel@biodiesel.org]; aweber@marciv.com[aweber@marciv.com]

Larry Schafer From:

Thur 2/27/2014 10:41:31 PM Sent:

Subject: Biodiesel Carbon Analysis Support Files

DESCRIPTION of Carbon Analysis Support Files Jan 2014.docx

ATT00001.txt

Jacobsen Feedstock Prices 2011 to 2013.xlsx

Overview of WAEES Model.pdf

Reassessment of life cycle greenhouse gas emissions for soybean biodiesel (4).pdf federal register notice (public document)

dgd-sum-120112.pdf

d1-15biodieselprofitability.xlsx

2010-3851.pdf

Biodiesel GHG Summary.xlsx

2014.01.15 FINAL VERSION OF CALCULATIONS.xlsx

15day-cornoil-bd-sum-022112.pdf

EIA Biodiesel Production Report Oct 2013 Data copy.pdf

Paul, Sharyn, David:

It was great meeting you and the rest of the EPA team last week. As discussed, attached are the support files for our analysis; apologies for the large size, but some of the PDFs are a bit big, and I thought you would find it helpful to have them all in one place. The shortcut/summary of our calculations is "2014.01.15 Final version of calculations.xls". Feel free to call if you have any questions or want to discuss further.

If you have questions or comments, then please let me know.

Thanks

Larry Schafer National Biodiesel Board O: 202.737.8801 M: 202.997.8072 LSchafer@Biodiesel.org

Biodiesel - America's Advanced Biofuel! www.americasadvancedbiofuel.com

1331 Pennsylvania Ave. NW Suite 505 Washington DC 20004

Carbon Analysis Support Files

Greenhouse Gas Emissions. These files provide the background support data that were utilized to calculate GHG reductions from various feedstock pathways (relative to petroleum diesel). The EPA final rule was the basis for most calculations. "Best available data" calculations included more recent work by USDA/University of Idaho, the CARB distillers corn oil pathway, and the CARB renewable diesel pathway. Files include the following:

- 2010-3851.pdf
- Reassessment of life cycle greenhouse gas emissions for soybean.pdf
- 15day-cornoil-bd-sum-022112.pdf
- dgd-sum-120112.pdf
- Biodiesel GHG Summary.xls.

Feedstock Utilization and Pricing. These files document the feedstocks utilized to produce biodiesel in the United States. The information from U.S. DOE-EIA is for the domestic biodiesel industry. There is no public information for feedstock utilization by renewable diesel producers. However, industry stakeholders provided input that only three (3) types of feedstock are currently utilized; animal fats, yellow grease, and distillers corn oil. Feedstock pricing information is from The Jacobsen, a fee based subscription service. Files include the following:

- EIA Biodiesel Production Report Oct 2013 Data copy.pdf
- Jacobsen Feedstock Prices 2011 to 2013

Biodiesel Production Costs. These files include the 3rd party production cost model from Iowa State University and a summary of both the GHG and cost calculations prepared by Bates White. In addition, an overview of the WAEES model utilized to forecast two scenarios in 2014 is available. Files include the following:

- D1-15biodieselprofitability.xlsx
- 2014.01.15 FINAL VERSION OF CALCULATIONS.xlsx
- Overview of WAEES Model. Pdf

	Soybean Oil (Illinois) crude/degu mmed	Canola Oil RBD (Chicago)	Bleachable Fancy - Tallow - Renderer (Chicago)	Choice White Grease (Chicago)	Yellow Grease (Illinois)	Stabiliazed Poultry Fat (Delmarva delivered)	Distiller's Corn Oil (IL)
2011 Jan	53.771	60.746	47.150	43.625	39.575	37.438	
Feb	54.137	61.032	47.513	45.974	41.395	43.645	
Mar	53.965	61.055	49.522	49.478	45.674	47.152	
Apr	56.639	63.171	51.588	50.850	46.300	48.375	
May	56.080	62.239	52.476	51.429	46.482	48.524	
Jun	55.729	61.626	54.048	52.546	47.028	45.227	
Jul	55.262	62.262	53.800	52.663	46.669	44.600	
Aug	54.350	61.595	49.565	49.348	44.707	45.304	
Sep	54.962	61.066	50.071	48.500	43.827	44.893	
Oct	51.992	59.011	47.600	46.351	43.202	44.393	
Nov	51.778	59.337	44.405	40.798	36.857	39.512	
Dec	50.292	58.209	47.119	41.202	36.015	37.708	
2012 Jan	51.108	58.133	44.200	40.300	35.125	38.200	37.313
Feb	52.406	59.281	45.750	45.265	38.850	40.863	43.375
Mar	53.420	61.056	48.045	49.000	42.182	45.750	44.818
Apr	54.942	62.948	47.300	47.389	41.038	45.188	42.175
May	50.771	60.385	48.932	49.131	41.625	44.784	42.477
Jun	49.552	60.367	45.548	44.803	37.869	41.952	37.762
Jul	54.419	64.308	45.452	43.756	36.810	40.012	36.476
Aug	52.329	63.731	44.630	44.810	37.174	40.283	37.174
Sep	53.433	64.228	45.526	45.197	38.132	41.026	39.763
Oct	50.187	62.070	39.717	38.000	34.690	37.913	36.065
Nov	48.317	59.417	34.000	33.903	30.500	35.100	29.988
Dec	49.617	58.736	36.300	36.727	33.238	36.450	34.500
2013 Jan	51.557	60.016	40.095	40.191	36.476	38.595	36.000
Feb	52.244	60.953	40.097	41.462	37.132	41.000	36.750
Mar	51.250	60.400	42.550	42.971	37.538	41.000	36.838
Apr	51.626	64.557	43.250	43.132	37.898	38.966	36.716
May	52.541	64.291	41.796	40.895	37.080	37.500	36.057
Jun	51.330	61.517	45.000	45.036	38.100	36.500	35.400
Jul	48.377	57.400	45.409	46.205	37.807	36.421	37.023
Aug	45.890	52.413	42.909	41.702	35.028	36.091	37.546
Sep	45.638	51.182	40.645	39.981	34.635	36.438	36.738
Oct	44.161	48.380	33.652	32.825	28.533	29.489	29.261
Nov	40.982	47.502	35.132	32.632	26.197	24.868	27.592
Dec	39.731	45.552	35.611	33.329	27.658	27.553	29.566

NOTE: The Jacobsen is a subscription based service. Information in this file cannot be utilized for other purp

must be cited.

Overview of WAEES Model

Prepared for:

Alan Weber

MARC IV

and

The National Biodiesel Board



Prepared by:

John Kruse, PhD

Director of Quantitative Analysis, WAEES

Email: jkruse@waees-llc.com

Phone: 573-228-9842


Website: www.waees-llc.com

1/19/2014

Overview of the WAEES Modeling System


The WAEES partial equilibrium modeling system is made up of a set of global econometric models emulating the behavior of the global agricultural sector. The partial equilibrium models can be broken down into crops, livestock and biofuels components encompassing feed grains, food grains, cotton, sugar, oilseeds, ethanol, biodiesel, beef, pork, and poultry.

WAEES Partial Equilibrium Models



The WAEES models cover 42 countries/regions with an additional 12 regional aggregates including the world total. WAEES follows USDA's reported data coverage which may mean that a zero is reported for a particular commodity which USDA does not cover or has discontinued covering. USDA currently covers at least 90 percent of global production; therefore, the countries which are omitted represent a small portion of total global production. Specifically the WAEES model includes Canada, Mexico, the United States, Caribbean and Central America, Argentina, Brazil, Bolivia, Chile, Paraguay, Uruguay, Other South America, the European Union 28, Other Europe, Russia, Ukraine, Uzbekistan, Other Former Soviet Union, Saudi Arabia, Turkey, Other Middle East, China, Japan, South Korea, Taiwan, Other East Asia, India, Pakistan, Other South Asia, Indonesia, Malaysia, Myanmar, Philippines, Thailand, Vietnam, Other Southeast Asia, Australia, Other Oceania, Egypt, Other North Africa, Kenya, South Africa, and Other Sub-Saharan Africa. WAEES also reports projections on crop area, yield and production for each of the EU-28 countries.

WAEES Regions follow the USDA Regions




Partial Equilibrium Models

Each partial equilibrium module is broken down into commodities with a system of structural equations capturing the supply and demand components for each of them. The drivers of these equations are theoretically derived based upon the behavioral postulates from economic theory of profit maximization by the market participants and utility maximization by consumers subject to various domestic and international trade policies. The diagram below illustrates the inter-linkages of the crops and livestock model. In the diagram, the blue boxes represent the key drivers (conditioning assumptions) of the agricultural sector including income, population, culture, inflation, exchange rates, domestic and trade policy, technology and input costs. The green boxes are an aggregate approximation of the crops sector. As relevant, each box represents an equation for each commodity covered. For example, there are specific feed demand equations for corn, sorghum, barley, soybean meal, sunflower meal, etc. The pink boxes are an aggregate approximation (within the diagram) of the detailed livestock sector model encompassing beef, pork and broilers. The diagram illustrates how income, population, and other factors drive food demand for crops and meats. Crude oil prices (and policies) drive the demand for biofuels. As demand increases, crop prices increase providing an incentive for production expansion. Technology growth drives yield expansion providing much of the needed production. Crop area may also grow to meet demand needs although in developed countries this often amounts to tradeoffs among crops. Ultimately supply and demand are balanced via commodity prices. If demand is stronger than supply, commodity prices increase until demand growth is slowed and supply growth is increases enough for supply and demand to balance.


Partial Equilibrium Modeling System

(Conceptual Framework Representation for One Country)



The WAEES partial equilibrium models solve iteratively to find equilibrium by balancing global supply and demand. This occurs at the individual country level for each commodity. Most countries are at least somewhat open to trade albeit with tariffs. The trade diagram below illustrates conceptually how global supply and demands are balanced within a "global" price equilibrium solution. Typically a large exporting country is chosen as the residual supplier for the world. The choice of this country does not affect the solution. The commodity price in the residual supplying country is solved for by assuming an initial level of exports. This price is then transferred to other countries through trade barriers, transportation costs, and exchange rates. Based on a given price level, each country determines how much it is willing to supply or demand at that price and subsequent how it wants to import or export. Occasionally a country has tariffs high enough that no trade will occur or only a fixed amount of trade will occur at the lower tariff level. Note that in those countries internal prices may not reflect the world level of prices because supply and demand must be balanced from domestic sources. After the supply and demand in each country is determined and the implied trade position, these trade positions are summed to find the new level of exports for the residual supplying country replacing the initial assumption. The process then repeats itself until prices adjust to balance global supply and demand. For example, if the sum of trade across all other countries is lower than the initial starting assumption for the residual supplying country, the price level in the residual supplying country will fall to balance supply and demand. This lower price level will then get transferred to all other countries affecting their supply and demand and ultimately net trade positions and of course replace the exports again in the residual supplying country. This process continues until global supply and demand balance.


How do partial equilibrium models solve for a global supply and equilibrium price?



An Example of the US Partial Equilibrium Model for the Biofuels Sector


Within the WAEES model, the US ethanol and biodiesel sectors are set up as partial equilibrium models with supply and demand equations and an endogenous ethanol and biodiesel price. The structure of the model has its roots in the ethanol specifications documented by John Kruse, Patrick Westhoff, Seth Meyer, and Wyatt Thompson in a 2007 journal article in AgBioForum entitled, "Economic impacts of not extending biofuel subsidies." With the second Renewable Fuel Standard, these original specifications have been updated to reflect the hierarchical system of mandates. The biofuels mandates require compliance with each specific mandate type including biodiesel, cellulosic, advanced and the overall renewable fuel mandate. The rational for different mandates in the legislation was to encourage biofuel producers to move towards feed stocks that provided the greatest level of greenhouse gas (GHG) reductions compared with conventional petroleum. The term "advanced biofuels" was used to describe biofuels that reduced GHG emissions by at least 50 percent compared with a 20 percent reduction requirement for conventional feed stocks. Cellulosic derived biofuels must reduce GHG emissions by 60 percent. Compliance with the mandates by the obligated parties is enforced by the EPA through as system of Renewal Identification Numbers (RINS) assigned to each type of biofuel produced. Obligated parties must demonstrated that they have met their assigned obligations through the number of RINS they have for each type of fuel. Theoretically there could be a specific RIN value for each type of mandate - cellulosic, biodiesel, advanced, and conventional, if each mandate was binding. Mandates are binding when the market is forced by policy to produce more than what normal economic conditions would suggest. The advanced biofuels are typically more expensive to produce than conventional biofuels resulting in those mandates being more binding than conventional biofuels mandates. Therefore RIN values (or prices) are typically significantly higher for advanced biofuels than conventional biofuels.

Hierarchical RINS Modeling




- Theoretically there can be 4 different RIN prices specific to each mandate if all the mandates are binding.
- Mandates are binding when the market is forced by policy to produce more than what normal economic conditions would suggest.
- Given the hierarchy of the mandates, it must be the case that RIN values for biodiesel are greater than or equal to advanced RIN values and advanced RIN values must be greater than or equal to conventional RINS. This is because biodiesel RINS can be used as advanced RINS and advanced RINS can be used as conventional RINS. (This process is referred to as demotion.)
- Biodiesel RINS can have the same value as advanced RINS if the biodiesel mandate is less binding than the advanced mandate.

US Biofuels Mandates in 2022



A detailed diagram of the US biofuels models is presented below. The demand for biofuels is largely mandate driven. However, if crude oil price edge higher it is possible for ethanol demand to be driven by market forces although the blend wall presents another hurtle. The supply of biofuels is driven by the profit margins of the biofuel plants. Profit margins are derived by subtracting the cost of the feed stocks and other variable costs of production from the valued of the products. In the case of ethanol, the value of the ethanol plus the value of the byproducts including corn oil and distiller's grains form the gross returns. The cost of ethanol is composed of the feed stock cost, primarily corn, and the other inputs. In the case of biodiesel, the value of biodiesel and the byproduct glycerin form the gross returns. The cost of producing biodiesel is composed of the feed stock costs such as vegetable oils, waste oils, corn oil and other inputs. The respective margins for ethanol and biodiesel drive capacity expansion in the longer term and capacity utilization in the short term for each sector. Equilibrium between biodiesel supply and demand is found by solving for the biodiesel price.

US Biofuels Partial Equilibrium Models



The WAEES Global Modeling Process

Forecast Assumptions

WAEES begins each semi-annual forecast by developing a set of conditioning assumptions that will be used for the forecast. These assumptions include the critical domestic and trade policies affecting agriculture and biofuels in each country; macroeconomic conditions such as per capita income growth, population growth, inflation rates, and exchange rates; technology assumptions such as crop yield growth; and key cost of production drivers such as interest rates, petroleum prices, wage rates, and other trends in tastes and preferences. Infrastructure constraints and land area expansion assumptions are also outlined in this process. These assumptions are direct inputs into the WAEES global agricultural partial equilibrium models.

Historical Data

The second step in the process is updating all historical data to the latest numbers. A large portion of the historical supply and demand data is drawn from USDA's Production, Supply, and Disposition (PSD) database. Historical data on crop area, yield, and production for each of the EU-28 countries is taken from Eurostat and supplemented with data from each of the country Ministries of Agriculture as needed. Some historical data such as sugarcane and sugar beet area harvested is taken from FAOSTAT, but the data is reviewed for consistency prior to being used in the models. Historical data on commodity prices are taken from a variety of sources including the respective Ministries of Agriculture (or equivalent) in each country, USDA, FAO, etc. Historical government policy information is gathered from USDA Gain Reports, the WTO, OECD, FAO, and the respective Ministries of Agriculture (or equivalent) in each country.

The timing of historical data releases determines when the WAEES forecasts are completed. The critical updates for PSD's global livestock data occur in April and October. The global crops data is updated more frequently throughout the year. Since the size of the southern hemisphere crops are generally available in April/May and the size of plantings in the northern hemisphere crops are generally known, WAEES conducts the first of the semiannual forecasts over the month of May targeting the beginning of June for release of the forecast numbers. The second forecast is typically done over the month of November targeting the beginning of December for a release of the forecast numbers. At this time of the year, the northern hemisphere crop sizes and the southern hemisphere plantings are generally known.

Model Development and Equation Updates

The WAEES global partial equilibrium models are in a constant state of review to ensure that the equation are performing adequately, the model structure is adapted to changes in the marketplace, changes in data sources are captured, and new coverage is added as necessary. While WAEES does not keep an exact count on the number of equations in the system, it now exceeds 20,000 equations. The performance of the behavioral equations within the system are continuously monitored within the system based on their percent root mean square errors, consistency with market behavior, and their recent pattern of historical errors. Prior to each forecast, the equations are reviewed and replaced as needed.

Model Calibration and Adjustment

After the historical data has been updated, each equation is recalibrated to the updated historical data. After reviewing the equation performance as per the description above, the model adjustment factors are set for the first forecast year. These adjustments are set based on a weighted average of the

equations errors over the previous 3-5 years in the model. In 99.5% of the equations this adjustment factor is held constant over the forecast horizon of 2013 through 2030. There are a few equations, particularly in the livestock sector, where adjustments are used to generate the livestock cycles.

Generating the forecast

After capturing the forecast assumptions, updating the historical data, reviewing the model equations, and calibrating the model, the model is then solved to generate a global forecast of commodity prices that balances supply and demand within each country and around the world. Since the commodities are highly interrelated within the model sometimes the forecast assumptions generate unexpected results and/or push the model into a region outside the experience based on historical data. The global solution is carefully reviewed and the equation results are evaluated based on direction and magnitude of response, and if necessary, the model equations are adjusted and the model is re-solved for a new global solution. These corrections are usually small or not needed, but some scenarios can push the model into untested ranges.

REASSESSMENT OF LIFE CYCLE GREENHOUSE GAS EMISSIONS FOR SOYBEAN BIODIESEL

A. Pradhan, D. S. Shrestha, J. Van Gerpen, A. McAloon, W. Yee, M. Haas, J. A. Duffield

ABSTRACT. This study updates the life cycle greenhouse gas (GHG) emissions for soybean biodiesel with revised system boundaries and the inclusion of indirect land use change using the most current set of agricultural data. The updated results showed that life cycle GHG emission from biodiesel use was reduced by 81.2% compared to 2005 baseline diesel, When the impacts of lime application and soil N2O emissions were excluded for more direct comparison with prior results published by the National Renewable Energy Laboratory (NREL), the reduction was 85.4%. This is a significant improvement over the 78.5% GHG reduction reported in the NREL study. Agricultural lime accounted for 50.6% of GHG from all agricultural inputs. Soil N₂O accounted for 18.0% of total agricultural emissions. The improvement in overall GHG reduction was primarily due to lower agricultural energy usage and improved soybean crushing facilities. This study found that soybean meal and oil price data from the past ten years had a significant positive correlation ($R^2 = 0.73$); hence, it is argued that soybean meal and oil are both responsible for indirect land use change from increased soybean demand. It is concluded that when there is a strong price correlation among co-products, system boundary expansion without a proper co-product allocation for indirect land use change produces erroneous results. When the emissions associated with predicted indirect land use change were allocated and incorporated using U.S. EPA model data, the GHG reduction for biodiesel was 76.4% lower than 2005 baseline diesel.

Keywords. Biodiesel, Biofuel, Greenhouse gas emissions, Land use change, Life cycle analysis, Soybean.

iofuels are becoming popular alternatives to fossil fuels, with state and federal policies, such as the Renewable Fuel Standard (RFS2), significantly increasing their demand over the past several years (EPA, 2010a). Although biofuels have the potential to become completely renewable, their production with today's technology requires some nonrenewable resources. e.g., synthetic fertilizers are used to improve yields, and fossil fuels are used for powering farm equipment.

The first comprehensive life cycle inventory (LCI) for biodiesel (BD) produced in the U.S. from soybean oil was published by the National Renewable Energy Laboratory (NREL) (Sheehan et al., 1998). The purpose of the NREL study was to conduct a life cycle assessment (LCA) to quantify the energy and emissions associated with the production and use of soybean biodiesel and compare it to petroleum diesel. The study took into account the emissions

by the USDA. The data for soybean crushing came from a performance study conducted in 1981. The study used a 1994 transesterification model from a single commercial transesterification facility. The NREL study reported that soybean biodiesel reduced carbon dioxide (CO2) emissions by 78.5% compared to petroleum diesel. The reason behind this reduction is that biomass-derived fuels participate in the relatively rapid cy-

associated with soybean agriculture, transport, crushing, oil

transesterification, biodiesel transport, and use of biodiesel

in a city bus. The study used 1990 soybean production data

from the Farm Costs and Return Survey (FCRS) conducted

cling of carbon to and from the atmosphere. Biomassderived carbon that ended up as CO2 leaving the tailpipe of a city bus was subtracted from the total CO2 as part of the biological recycling of carbon.

The objective of this study is to update the life cycle greenhouse gas (GHG) emissions calculations based on the most recent complete set of data for soybean biodiesel production via base catalyzed transesterification. Even though more current partial data for agriculture were available, the data used in this article are from 2006. It is important to use agricultural data from a single year, as agricultural data vary significantly from year to year depending on factors such as weather and pest infestation. The year 2006 was the most recent year that had a complete set of agriculture data available. This study compares the new LCA result with the NREL result and provides an explanation of the reasons for any differences. This study also points out the potential pitfalls of the system boundary expansion approach for impact assessment of indirect land use change, including the

Submitted for review in December 2011 as manuscript number FPE 9548; approved for publication by the Food & Process Engineering Institute Division of ASABE in October 2012

The authors are Anup Pradhan, ASABE Member, Agriculture Engineer, Dev S. Shrestha, ASABE Member, Associate Professor, and Jon H. Van Gerpen, ASABE Member, Professor, Department of Biological and Agricultural Engineering, University of Idaho, Moscow, Idaho; Andrew J. McAloon, Cost Engineer, Winnie Yee, Chemical Engineer, and Michael J. Haas, Chemist and Lead Scientist, USDA-ARS Eastern Regional Research Center, Wyndmoor, Pennsylvania; and James Duffield, Senior Agricultural Economist, USDA Office of the Energy Policy and New Uses, Washington, D.C. Corresponding author: Dev S. Shrestha, Department of Biological and Agricultural Engineering, University of Idaho, P.O. Box 442060, Moscow, ID 83844; phone: 208-885-7545; e-mail: devs@uidaho. edu.

Transactions of the ASABE

Vol. 55(6): 2257-2264

2012 American Society of Agricultural and Biological Engineers ISSN 2151-0032

assumption that this approach will automatically account for co-product allocation.

METHODOLOGY

This study takes two different approaches; the first is the "base case," the methodology of which is consistent with the NREL's attributional LCA. Attributional LCA (ALCA) is a "business as usual" method that accounts for environmentally relevant physical flows to and from a product system. ALCA uses average values based on normal, current business practices. ALCA does not include any indirect effects that are not directly related to the production of biodiesel. The second approach is the "consequential" LCA, which includes factors such as indirect land use change. Consequential LCA (CLCA) aims to predict the consequences if changes are made to an established process. CLCA includes indirect changes in addition to direct effects.

The system boundary for the base case in this study is similar to that of the NREL study, except for inclusion of the use of agricultural lime (to improve soil pH) and soil ni-

T

trous oxide emission, and exclusion of oil transport. The list of inputs and outputs for the attributional LCA is shown in table 1. Agricultural lime was included in the base case because it is used periodically on soybeans and was inadvertently omitted from the NREL analysis. The impact of soybean oil transport was studied separately and not included in the base case because, for the most part, the soybean oil biodiesel plants considered in this study are colocated with soybean crushing plants. The GHG emissions were estimated from energy and material inputs in the production process. The emissions were calculated by multiplying the inputs by the corresponding emission factor. The data for estimating the effect of indirect land use change were borrowed from a recent EPA analysis (EPA, 2010b).

Three major anthropogenic greenhouse gases (CO₂, CH₄, and N₂O) were used to estimate the net GHG emissions. All emissions were reported as CO₂-equivalent (CO₂e) emissions. The CO₂e value indicates a GHG's global warming potential (GWP), as advocated by the Intergovernmental Panel on Climate Change (IPCC). GWP indicates the relative strength of radiative forcing (RF) of a GHG compared to CO₂ integrated over time. Therefore, CO₂ has a GWP of unity. The IPCC Second Annual Report

Inputs	Quantity Used	Embedded Energy	gCO ₂ e Factor	gCO ₂ e [e]
		Embedded Energy	gCO ₂ e Factor	gCO ₂ e · ·
Soybean agriculture	(per ha)	24.024.1.1.01	on marrial (c)	
Diesel	33.3 L [a]	35.9 MJ L ^{-1 [h]}	89.7 MJ ^{-1 [c]}	107,233.7
Gasoline	12.8 L [a]	32.4 MJ L-1 [h]	90.9 MJ ⁻¹ [c]	37,698.0
LP Gas	2.0 L [n]	23.7 MJ L-1 [b]	76.1 MJ ^{-1 (d)}	3,607.1
Natural gas	4.1 m ^{3 (a)}	36.6 MJ m ^{-3 [b]}	72.4 MJ ^{-1 [d]}	10,864.3
Nitrogen	3.3 kg [a]	58	3.6 g ^{-1 [d]}	11,880.0
Phosphorus	12.1 kg [*]	€8	1.2 g ^{-1 [d]}	14,520.0
Potassium	22.4 kg [*]	-80	0.8 g ^{-1 [d]}	17,920.0
Lime	463.7 kg [a]	#	0.6 g ⁻¹ [b]	278,220.0
Seed	68.9 kg [a]	. 40	189.3 kg ^{-1 [k]}	13,042.8
Herbicide	1.6 kg ^(f)		25.8 g-1 [d]	41,280.0
Insecticide	0.04 kg ¹⁷		30.0 g ^{-1 [d]}	1,200.0
Electricity	17.1 kWh [*]	3.6 MJ kWh ^{-1 [g]}	208.4 MJ ^{-1 [d]}	12,829.1
		15	Subtotal:	550,295.0
Soil N ₂ O emission	(per ha)			120,468.5
Soybean transport	(per ha)			56,464.3
Soybean crushing	(per L of BD)			
Electricity	212.3 Wh [h]	3.6 MJ kWh ^{-1 [g]}	208.4 MJ ^{-1 [d]}	159.3
Natural gas	0.11 m ^{3 [h]}	36.6 MJ m ⁻³ [b]	72.4 MJ ⁻¹ [d]	291.5
Hexane	11.1 g [b]	10 - 00	0.2 g ^{-1 [b]}	2.2
W	572		Subtotal:	453.0
Biodiesel conversion	(per L of BD)	304	57	
Electricity	44.6 Wh [b]	3.6 MJ kWh ^{-1 [g]}	208,4 MJ ^{-1 [d]}	33.4
Steam at 10.3 bar (150 psi)	124.1 g [h]	2.0 MJ kg ^{-1-[i]}	119.1 MJ ^{-1 lji}	29.6
Methanol	96.7 g [h]	20.1 MJ kg-1 [d]	67.7 MJ ^{-1 [d]}	131.6
Sodium methylate	2.7 g [h]		7.9 g ^{-1 (k)}	21.3
Hydrochloric acid	0.5 g [h]	920	13.5 g ^{-1 [k]}	6.8
			Subtotal:	222.7
Biodiesel transport and distribution	(per L of BD)			22.5
Biodiesel combustion	(per L of BD)			21.7

[[]a] 2006 ARMS and ERS data (data were obtained from the USDA through special request).

[[]h] ANL, 2010.

[[]c] DOE, 2008.

[[]d] EPA, 2010d.

Product of columns 2, 3 (when applicable), 4, and proper unit conversion factor to get emission in gCO₂e.

^[f] NASS, 2007.

Direct unit conversion.

[[]h] ARS model.

^[i] Steam table data

Natural gas as fuel with 60.8% boiler efficiency (steam generation at 150 psig = 1411 Btu lb⁻¹, and the enthalpy of evaporation from the steam table = 858 Btu lb⁻¹, which gives the total natural gas to steam usage efficiency of 858/1411 = 60.8%).

[[]k] Sheehan et al., 1998.

(SAR) assesses the GWP of CH₄ as 21 and N₂O as 310 for a 100-year horizon (IPCC, 1996). The IPCC Third Assessment Report (TAR) re-evaluates the GWP of CH₄ as 23 and N₂O as 296 for the same time horizon (IPCC, 2001). The United Nations Framework Convention on Climate Change reporting guidelines for national inventories were updated in 2006 but continue to require the use of GWP values from the IPCC SAR (UNFCCC, 2006). This requirement of using SAR GWP values ensures that new estimates of aggregate GHG emissions are consistent with estimates developed prior to the publication of the IPCC TAR and the IPCC Fourth Assessment Report (AR4), which re-evaluates the GWP of CH₄ as 25 and N₂O as 298 (IPCC, 2007). In order to comply with UNFCCC reporting standards, this article uses SAR GWP values. The U.S. EPA also follows UNFCCC guideline and uses GWP values from SAR (EPA, 2010c) in its renewable fuel standard (RFS2) life cycle analysis.

The GHG emissions for soybean biodiesel production were expressed as grams of CO₂-equivalent (gCO₂e). The energy inputs were multiplied by the embedded energy (low heating value for all fossil fuels) of the input and then multiplied by the appropriate GHG factor (table 1). For the non-energy inputs, where energy equivalence is not applicable, the input was directly multiplied by the GHG factor to calculate gCO₂e. The results were compared with 2005 baseline diesel GHG emissions, as required by The Energy Independence and Security Act (EISA) of 2007, to quantify the relative benefits of soybean biodiesel.

DATA DESCRIPTION AND ASSUMPTIONS Soybean Agriculture

At the time of the Sheehan et al. (1998) study, the most recent soybean production data were from the USDA 1990 Farm Costs and Return Survey (FCRS). In this article, all farm input and direct energy data for soybean production are from 2006, the most recent set of soybean survey data available at the time of this study. Agricultural inputs and outputs, such as yields and use of pesticides, vary from year to year. Therefore, mixing and matching agricultural data from different years can produce an unrealistic picture. Temporal variation could be minimized by averaging several years of data, but complete sets of agricultural data are not generally available for multiple consecutive years. Therefore, a complete set of the most recent agriculture data from a single year was used in this study.

In order to ensure that 2006 was not an abnormal year, which could bias the result, we carried out a linear regression analysis on yields from 1980 to 2010. This analysis verified that the yield for 2006 was within the 95% confidence interval (36.4 to 48.8 bu ac¹) of predicted yield. The fertilizer, lime use, and direct energy use (such as diesel, gasoline, and natural gas consumption) were from the 2006 Agricultural Resource Management Survey (ARMS) and National Agricultural Statistics Service (NASS) data compiled by the USDA Economic Research Service (ERS). Chemical data for 2006 were from a chemical survey conducted by NASS (NASS, 2007). The 2006 ARMS and NASS soybean survey provided detailed state-level data for 19 major U.S. states. The state soybean yield data were es-

timates reported by NASS (NASS, 2010). The national average yield was 2906.7 kg ha⁻¹ (43.2 bu ac⁻¹) in 2006. The soybean farm survey data were weighted by state acreage to derive the average quantity used for U.S. soybean production. The CO₂e emission values were from the Excel sheet "emission factors" in EPA data (EPA, 2010d). The CO₂, CH₄, and N₂O emissions for hexane and agricultural lime, not provided in the EPA report, were from the Excel sheet "BD" in the GREET (Greenhouse gases, Regulated Emissions, and Energy use in Transportation) model (ANL, 2010). These CO₂e emission values were then converted to CO₂e factors using SAR GWP values.

Soil Emission Data

This study used soil N₂O emissions (not available at the time of the NREL study) from the GREET model (ANL, 2010). N₂O is emitted through (1) direct emissions (including nitrification, denitrification, and volatilization) from the soil to the air, and (2) indirect emissions (including leaching and runoff of nitrate into waters) (Huo et al., 2009). N₂O emissions from the biological fixation of nitrogen are not included in the model, as the IPCC, in the 2006 guidelines, removed biological fixation of nitrogen as a direct source of N₂O (IPCC, 2006).

The GREET model (ANL, 2010) estimates soil N₂O emission using the total amount of nitrogen in the soybean biomass left in soybean fields (aboveground and belowground biomass) and in the nitrogen fertilizer applied. GREET estimates 7.4 g of nitrogen in the biomass per kilogram of soybean produced (200.7 g N bu⁻¹ soybean). IPCC suggests an average conversion factor of 1% for the production of N₂O from biomass nitrogen and fertilizer nitrogen (IPCC, 2006). To estimate the total N₂O emission, 1% of the summed nitrogen content from biomass and synthetic fertilizer was multiplied by the factor 1.57 to account for the ratio of the molecular weights of N₂O and N₂, per IPCC recommendations. Using values from the GREET model, the N₂O emission from the soil was estimated to be 388.1 g N₂O ha⁻¹ (3.63 g N₂O bu⁻¹).

Soybean Transport

The average hauling distance for soybeans from the point of production to that of processing depends on the crushing capacity of the plant. For an oil crushing plant with an annual capacity of 378 million L (100 million gal), the theoretical minimum hauling distance was calculated to be 56 km (35 mi), assuming a corn-soybean rotation and that the crushing plant was located at the center of a squareshaped agricultural area from which it draws the soybeans (Biodiesel Education, 2012). Because of system inefficiencies, the actual hauling distance would be greater than this. A one-way trip of 81 km (50 mi) was assumed to be the average distance to haul soybeans to the crushing/biodiesel plant using a truck as the mode of transportation (ANL, 2010). This estimation was based on 16 km (10 mi) to transport soybeans from farm to storage and another 64 km (40 mi) to transport soybeans to the crushing/biodiesel plant. The GREET model estimates of CO2, CH4, and N2O emissions for soybean transportation were 512.32, 0.5886, and 0.0133 g bu⁻¹, respectively. Using SAR GWP, the CO₂e for soybean transport was estimated to be 56,464.3 g ha

(529 g bu⁻¹) (table 1). The theoretical analysis provided a means of data verification.

Soybean Oil Extraction and Transesterification

This study uses the energy input data for soybean crushing, hexane extraction of the oil, and biodiesel production via alkali-catalyzed transesterification from a biodiesel plant model developed by the USDA-ARS using SuperPro designer (Intelligen, Inc., Scotch Plains, N.J.). The ARS model was prepared from process designs, equipment specifications, costs, and energy consumptions that were provided by technical experts and equipment suppliers to the soybean crushing and biodiesel industry. The model estimates the electrical and thermal energy inputs required for hexane extraction and its subsequent refining and conversion to biodiesel at an annual scale of 38.6 million L (10.2 million gal) of biodiesel, 137,491 Mg of soybean meal, 8,167 Mg of soybean hulls, and 3,975 Mg of crude glycerin. The model used in the analysis allows the plant to generate its own steam from natural gas with a life cycle efficiency of 60.8% (table 1). The model does not represent an industry average, but it provides a blueprint of a specific biodiesel plant based on the best information available from equipment manufacturers and communication with the industry.

Biodiesel Transport, Distribution, and Combustion

The biodiesel transport and distribution data used in this study were taken from the GREET model (ANL, 2010), which estimates a one-way trip of 540 km (335 mi) for biodiesel transport and distribution using a combination of truck, barge, and rail. This estimation was based on 52 km (32 mi) by truck, 68 km (42 mi) by barge, and 373 km (232 mi) by rail to transport the biodiesel to a distribution center, and another 48 km (30 mi) by truck to transport the biodiesel to its final destination. The GREET estimates of CO₂, CH₄, and N₂O emissions for biodiesel transport and distribution were 704.7, 0.81, and 0.0167 g mmBtu⁻¹ BD, respectively. Using SAR GWP, the CO₂e for biodiesel transport and distribution was estimated to be 0.7 g MJ⁻¹ fuel (22.5 g L⁻¹ BD) (table 1).

The CO₂ emission from biodiesel combustion was not included in the model because it is assumed to be equal to the amount of CO₂ captured by soybeans during photosynthesis. Exclusion of CO₂ emission is consistent with the NREL study. The combined N₂O and CH₄ emission from biodiesel combustion was estimated to be 21.7 gCO₂e L⁻¹ BD (EPA, 2010d) (table 1).

CO-PRODUCT ALLOCATION

In order to provide a consistent comparison to the NREL report, this study used a mass-based allocation method that allocates energy and emissions to the various co-products by their relative weights. The USDA Economic Research Service (ERS, 2009) reported a 2006-2007 U.S. average oil yield of 0.189 kg oil kg⁻¹ soybean (11.34 lb bu⁻¹). This extraction rate is higher than the 0.169 kg oil kg⁻¹ soybean (10.16 lb bu⁻¹) used in the NREL study. The oil extraction rate for crop year 2006-2007 was used in this study in order to be consistent with the 2006 ARMS agricultural input da-

Table 2. Base case emissions for biodiesel with co-product allocation.

	Allocation	Emiss (gCO ₂ e GJ ⁻¹	
Subsystem	Factor (%)	Before Allocation ^[a]	After Allocation
Soybean agriculture	18.4	28,128.9	5,175.7
Soil N2O emission	18.4	6,157.9	1,133.1
Soybean transport	18.4	2,886.2	531.1
Oil recovery	18.4	13,853.2	2,549.0
Biodiesel conversion	89.9	6,810.4	6,122.5
Biodiesel transport	100	688.1	688.1
Biodiesel combustion	100	663.1	663.1
Total		59,187.8	16,862.6
Diesel emissions (gCO ₂ e GJ ⁻¹ diesel) ^[b]			89,668.2
GHG reduction for biodiesel			81.2

From table I (last column of table I was converted to gCO₂e GJ⁻¹ biodiesel using conversion factors of 19,563.3 MJ of energy from biodiesel ha⁻¹ and 32.7 MJ L⁻¹ of biodiesel).

ta. After excluding the hulls and waste material, the soybean produced 20.5% oil and 79.5% meal by weight. Total emissions from biodiesel were allocated between oil and meal accordingly.

Transesterification of soybean oil produces biodiesel and crude glycerin. The NREL model of transesterification used a biodiesel to crude glycerin production ratio of 4.7:1 (10,504 kg h⁻¹ for biodiesel and 2,235 kg h⁻¹ for crude glycerin). According to this ratio, the NREL study allocated 82.4% to biodiesel and 17.6% to crude glycerin. However, modern plants have biodiesel to crude glycerin ratios of about 10:1 by weight (da Silva et al., 2009; Thompson and He, 2006; Van Gerpen et al., 2006). The model in this study uses output rates of 4,256.3 kg h⁻¹ for biodiesel and 479 kg h⁻¹ for crude glycerin. This corresponds to a ratio of 89.9% biodiesel to 10.1% crude glycerin, which is close to the modern industrial average. The co-product share of crude glycerin was deducted from the estimated GHG emissions of soybean agriculture, soybean transport, and oil recovery. The overall allocation for soybean agriculture, soybean transport, and oil recovery was therefore 18.4% (20.5% × 89.9%), as shown in table 2.

RESULTS AND DISCUSSION

The average soybean yield was 2,907 kg ha⁻¹ (43.2 bu ac⁻¹) in 2006 (NASS, 2010). With 0.189 kg oil kg⁻¹ soybean and 96% conversion efficiency from oil to biodiesel by weight, each hectare of soybean production is equivalent to 598.7 L of biodiesel (64.12 gal BD ac⁻¹). Biodiesel has a lower heating value (LHV) of 32.7 MJ L⁻¹ (Sheehan et al., 1998).

The gCO₂e values from table 1 were converted to consistent units of gCO₂e per GJ of biodiesel output (table 2). The conversion used was 1 ha of soybean production is equivalent to 19,563.3 MJ of energy from biodiesel.

The reduction in GHG emission (81.2%) compared to the reduction reported by NREL (78.5%) was mainly because of improved agricultural management practices and increased energy efficiency in soybean crushing. Since the time of the NREL study, soybean yield has consistently im-

[[]b] DOE, 2008.

proved due to genetically engineered varieties, improved chemical applications, and new management practices (Ash et al., 2006). For example, in conjunction with reduced chemical applications and improvements in management practices, fewer equipment trips across the fields are required. Largely as a result of this, diesel fuel use decreased from 49.4 L ha-1 (5.29 gal ac-1) in 1990 to 33.3 L ha-1 in 2006, and gasoline use decreased from 29.0 to 12.8 L ha⁻¹ during the same period. In addition, recently constructed soybean crushing facilities are more energy efficient than older facilities. For instance, since 2002, the U.S. EPA has required soybean plants to limit their hexane use (EPA, 2001). Currently acceptable levels of hexane loss are less than one-third of the level reported in the NREL study (Woerfel, 1995). As a consequence, the new hexane input value used in this study is one-half of that reported in the NREL study.

EFFECT OF ADDING AGRICULTURAL LIME

The NREL study did not consider the impact of agricultural lime usage on GHG production. Lime is added periodically to reduce soil acidity and to increase soybean yield. The average lime application for soybean production for crop year 2006 was 463.7 kg ha⁻¹ (NASS, 2007). With 0.6 gCO₂e g⁻¹ applied lime (CaCO₃) (ANL, 2010), the CO₂ emission associated with lime use was estimated to be 278,220.0 gCO₂e ha⁻¹. The GHG emission for lime was mainly from mining and processing. Of all agricultural inputs, lime contributed the most GHG. In fact, the emission from lime was 50.6% of the total GHG from agriculture inputs and 2.5 times more than the emission from diesel use. the next largest source of GHG emissions from agricultural inputs. Therefore, lime adds a significant amount of GHG emission to the soybean biodiesel life cycle assessment. The inclusion of lime was also recommended by Landis et al. (2007). The main reason for this high emission from lime is that the quantity of lime applied is significantly higher than other inputs (table 1). If agricultural lime was not included, for a more direct comparison to the NREL report, then the GHG reduction from the use of biodiesel relative to petroleum diesel would have been 84.1%, compared to the 81.2% value in table 2.

EFFECT OF ADDING N2O EMISSIONS FROM SOILS

The NREL estimate did not include soil N_2O emissions. N_2O emissions accounted for 18.0% of total GHG emission from soybean agriculture (emission from agricultural inputs plus soil N_2O). If soil N_2O emissions were not included in the base case study, then the reduction in GHG emission from the use of biodiesel would have been 82.5% rather than 81.2%. The soil N_2O emissions contribute 6.7% of total life cycle GHG emissions for biodiesel production, and hence cannot be neglected. If both lime and N_2O emission were excluded from the life cycle inventory, for a direct comparison with the NREL results, then the GHG reduction for biodiesel relative to petroleum diesel would have been 85.4%, compared to 78.5% reported in the NREL study.

EFFECT OF ADDING SOYBEAN OIL TRANSPORT

The base case estimation did not include emissions associated with soybean oil transport because this study assumed that the soybean crushing facility and biodiesel conversion plant were co-located. However, several biodiesel plants purchase oil and transport it to their plant. The NREL study included oil transport in its life cycle inventory, which added 560.9 gCO₂e GJ⁻¹ BD for 919 km (571 mi) using rail as a mode of transportation. This is equivalent to 61.0 gCO₂e GJ⁻¹ BD for 100 km of oil transport. This is only 1% of the emissions for biodiesel conversion. Thus, if oil transport from the crushing site to the biodiesel production site is a short distance, then emissions from oil transport can be neglected without causing much error.

EFFECT OF LAND USE CHANGE (LUC)

In addition to direct emissions, the 2007 EISA requires that calculations of life cycle GHG emissions include all significant indirect emissions, such as significant emissions from indirect land use changes (ILUC). The LUC estimates (both direct and indirect) used by the EPA include domestic and international land use conversions induced by increased consumption of renewable fuels in the U.S. A summary of the EPA calculations for GHG emissions from LUC is shown in table 3.

International land use change is land use change in all countries other than the U.S. How land is used is assumed to be determined by the relative profits from various activities. The EPA estimated the land use change impact with a 30-year horizon beyond the year 2022, when RFS2 is fully implemented, with a 0% discount rate for its rulemaking (EPA, 2010c). A 0% discount rate means that the GHG emissions today are worth the same as emissions 30 years from now. To calculate the annual land use change impact for the next 30 years, an emission average was calculated using the following equation:

$$LUC_GHG = \frac{\sum_{n=0}^{29} LUC_n}{30}$$
 (1)

where LUC_GHG is the annual GHG per GJ of biodiesel, and LUC_n is the GHG emission due to land use change in the nth year. Year 0 in equation 1 is the year 2022. The LUC GHG value estimated from this equation using data

Table 3. Summary calculation of annual life cycle GHG emission from LUC for the year 2022 and beyond (Source: EPA, 2010b).

	Emission (gCO ₂ e GJ ⁻¹ Biodiesel) ^[n]		
Emission Category	Year 0	Years 1-19	Years 20-99
International land use change	1,114,419	5,078	-114
Domestic soil carbon ^[h]	-252,977	0	0
Domestic livestock	-1,991	-1,991	-1,991
Domestic rice methane	-7,536	-7,536	-7,536
International farm inputs and fert. N2O	5,120	5,120	5,120
International livestock	-6,100	-6,100	-6,100
International rice methane	2,066	2,066	2,066
Total	853,001	-3,363	-8,555

The conversion factor 1 GJ = 0.948 mmBtu was used to convert to gCO₂e GJ⁺ biodiesel from the original EPA calculations.
Average domestic soil carbon was used for years 1-19 and 20-99.

55(6): 2257-2264

from table 3 is 23,452 gCO₂e GJ⁻¹ biodiesel. The EPA assumed that LUC_GHG in equation 1 is caused solely by the shift in the equilibrium of demand for soybean oil and that no allocation of land use change GHG to the co-products is needed (EPA, 2010e). In other words, increased demand for oil is the only driving force in shifting the equilibrium. This assumption is based on the economic principle that assumes when oil price goes up, more soybeans will be crushed, thus increasing the oil supply, and as a result, the supply of meal also increases. With a static demand for meal, as supply increases, the meal price would go down, in which case, it could be argued that the meal is not a driving factor in LUC. Contrary to this assumption, the recent price trend data for soybean meal and oil show that they both go up simultaneously.

The USDA-ERS price data for oil and meal over the past ten years were regressed. The prices of oil and meal had a statistically significant positive correlation, with R^2 = 0.73 (p < 0.0018) (fig. 1). Another source (IndexMundi, 2012) of historical monthly commodity price data showed that soybean oil and soybean meal prices had a positive correlation, with $R^2 = 0.80$ (p < 0.0001) for the period 2002 to 2012. Soybean prices also increased during the same period. The soybean meal price could have increased just because of the higher soybean prices, and the correlation we observed could thus have been just an artifact of an increasing soybean price. To test if this was the case, a relative increase in oil price was compared to a relative increase in meal price. It is important to compare the relative increases in price, as oil has a much higher price per unit of mass compared to meal. For a relative comparison, the prices of oil, meal, and soybean were normalized using equation 2:

Equation 2 linearly scales the prices of oil, meal, and soybean between 0 and 1. The normalized prices for oil and meal were regressed with the normalized price of soybeans




Figure 1. Annual average price of soybean oil and meal from 2000 to 2009 (source: ERS, 2011).

using equations 3 and 4:

Normalized oil price =
$$a_1 + b_1 \times \text{Normalized soybean price}$$
 (3)

Normalized meal price =
$$a_2 + b_2 \times \text{Normalized soybean price}$$
 (4)

The slopes of the linear lines (b_1 and b_2) represent the relative increase in the price of oil or meal compared to the price of soybeans. The regression analysis showed $b_1 = 0.94$ and $b_2 = 1.00$. This result tells us that the oil price increased by only 94 cents per dollar increase in the soybean price, whereas the meal price increased dollar per dollar with the soybean price. Since the slope of the normalized meal price (b_2) was greater than the slope of the normalized oil price, it was concluded that the relative price of meal was increasing at least as rapidly as the price of oil.

A strong positive correlation between meal and oil indicates that demand for meal and oil increase proportionally. From these results, it was concluded that the price of soybean meal is as strong an incentive to trigger LUC as soybean oil. If price is the driving force, then both meal and oil are the drivers for LUC. In the EPA analysis, the system was expanded to include soybean meal in the partial equilibrium model, which assumes constant meal use. The assumption of constant meal use effectively allocates all LUC emissions to soybean oil, as the model assumes that soybean oil is the only driving factor for LUC.

The strong positive correlation between oil and meal price shows that both co-products act together as a unified driving force in any resulting LUC impact. The extent to which meal should be held accountable for indirect land use change depends on the correlation between meal and oil prices. If there were no positive correlation, then oil price increases alone could be blamed for all indirect land use change, and all LUC_GHG could be attributed to oil, as was done in the EPA study (EPA, 2010d). However, since there is a statistically significant positive correlation between oil and meal prices, the LUC_GHG effects should be allocated to both meal and oil. Thus, equation 1 becomes:

$$LUC_GHG = \frac{n=0}{30} \times Allocation factor$$
 (5)

where the allocation factor partitions the GHG impact to its meal and oil sources. This equation takes into account the fact that both soybean meal and oil are responsible for LUC_GHG, and it attempts to identify the proportion of this value that is attributable to soybean oil production. Assuming the same soybean oil allocation factor for indirect land use change that was applied for soybean agriculture (18.4%), the LUC_GHG was estimated to be 4,315 gCO₂e GJ⁻¹ BD (instead of the value of 23,452 gCO₂e GJ⁻¹ in the absence of allocation).

The reduction in GHG emissions from the use of biodiesel, compared to 2005 baseline diesel, was 76.4% after inclusion of LUC (compared to 81.2% before inclusion). The GHG reduction of 76.4% was significantly greater than

2262 TRANSACTIONS OF THE ASABE

the 57% reduction reported by the EPA in its RFS2 rule-making (EPA, 2010c). The difference arises from the application of the allocation factor to partition GHG impact between oil and meal. The EPA report assigned an allocation factor of 100% to soybean oil and hence to biodiesel. That is, the total GHG impact of land use change was attributed only to biofuel. If a similar assumption is made in this study, then the GHG reduction is estimated to be 55.0% which is close to the value of 57% that was reported by the EPA.

SUMMARY AND CONCLUSIONS

Using the most recent set of agricultural data available, from the 2006 crop year, soybean biodiesel production and usage were calculated to result in an 81.2% reduction of GHG emissions relative to those calculated for petroleum diesel usage based on 2005 data. This calculation incorporated agricultural lime application and soil N₂O emissions. If lime and N₂O were not included, for a more direct comparison with the 1998 NREL study, the reduction would have been 85.4%. This is a significant improvement over the 1998 NREL study, which reported a total GHG reduction of 78.5%. The improvement in GHG emission reductions was mainly due to reduced agricultural energy usage and improved energy efficiency in modern soybean crushing facilities.

The base case in this study used a similar system boundary as the 1998 NREL study, except that agricultural lime use and soil N2O emissions were added, and soybean oil transport was omitted. Lime contributed about 50% of the total GHG emissions from soybean agriculture. The GHG emission from lime use was about 2.5 times higher than that of diesel use, the second highest contributor of GHG emissions from agricultural inputs. The emission from soil N2O was about 18.0% of total emissions from agricultural and 6.7% of the total biodiesel life cycle GHG emissions. Therefore, it was concluded that soil N2O emissions are significant and cannot be neglected. The impacts of oil transport were excluded from the base case in this study because most biodiesel plants are co-located with oil crushing facilities. The analysis revealed that GHG emission from oil transport of 100 km was equivalent to only 1% of the GHG emission from transesterification. Therefore, the GHG emission from oil transport for short distances could be neglected without causing much error in the final result.

Because soybean oil prices had a strong positive correlation with meal prices, it was argued that both meal and oil prices are responsible for shifting the equilibrium of soybean demand. Holding only soybean oil responsible for land use changes, with the assumption that soybean meal price does not change or decrease because of increased meal supply, was found to be erroneous. When the emissions associated with land use change (direct and indirect) were incorporated into the base case results, the net GHG reduction from biodiesel use was found to be 76.4% less than the emissions for 2005 baseline diesel.

REFERENCES

- ANL. 2010. The greenhouse gases, regulated emissions, and energy use in transportation (GREET) model V1.8d.1.

 Argonne, Ill.: U.S. DOE Argonne National Laboratory.

 Available at: http://greet.es.anl.gov/. Accessed 14 June 2011.
- Ash, M., J. Livezey, and E. Dohlman. 2006. Soybean backgrounder. OCS-2006-01. Washington, D.C.: USDA Economic Research Service. Available at: www.ers.usda.gov/ publications/OCS/apr06/OCS200601/OCS200601_lowres.pdf. Accessed 14 June 2011.
- Biodiesel Education, 2012. Minimum hauling distance calculator. Moscow, Idaho: University of Idaho. Available at: www.cals.uidaho.edu/bioenergy/calculators.asp. Accessed 10 April 2012.
- da Silva, G. P., M. Mack, and J. Contiero. 2009. Glycerol: A promising and abundant carbon source for industrial microbiology. *Biotech. Advances* 27(1): 30-39.
- DOE. 2008. Development of baseline data and analysis of life cycle greenhouse gas emission of petroleum-based fuels. DOE/NETL-2009/1346. Morgantown, W.V.: DOE National Energy Technology Laboratory. Available at: www.netl.doe. gov/energy-analyses/pubs/NETL%20LCA%20Petroleum-Based%20Fuels%20Nov%202008.pdf. Accessed 14 June 2011.
- EPA. 2001. National emission standards for hazardous air pollutants: Solvent extraction for vegetable oil production: Final rule. Federal Register 66(71):19006-19026.
- EPA, 2010a. EPA finalizes regulations for the National Renewable Fuel Standard Program for 2010 and beyond. EPA-420-F-10-007. Washington, D.C.: U.S. Environmental Protection Agency. Available at: www.epa.gov/otaq/renewablefuels/ 420f10007.pdf. Accessed 14 June 2011.
- EPA. 2010b. Biodiesel from soybean oil and waste grease. EPA-HQ-OAR-2005-0161-3173.9. Washington, D.C.: U.S. Environmental Protection Agency. Available at: www.regulations.gov/#!documentDetail;D=EPA-HQ-OAR-2005-0161-3173.9. Accessed 14 June 2011.
- EPA. 2010c. Renewable Fuel Standard Program (RFS2) Regulatory Impact Analysis. EPA-420-R-10-006. Washington, D.C.: U.S. Environmental Protection Agency. Available at: www.epa.gov/otaq/renewablefuels/420r10006.pdf. Accessed 14 June 2011.
- EPA. 2010d. Lifecycle greenhouse gas (GHG) impacts calculations. EPA-HQ-OAR-2005-0161-3174.10. Washington, D.C.: U.S. Environmental Protection Agency. Available at: www.regulations.gov/#!documentDetail;D=EPA-HQ-OAR-2005-0161-3174.10.
- EPA. 2010c. Lifecycle GHG analysis. In Renewable Fuel Standard Program (RFS2) Regulatory Impact Analysis (EPA-420-R-10-006), 298-502. Washington, D.C.: U.S. Environmental Protection Agency. Available at: www.epa.gov/otaq/renewablefuels/420r10006.pdf.
- ERS. 2009. Table 09: Soybeans: Monthly value of products per bushel of soybeans processed, and spot price spread, U.S., 1990/91-2007/08. In *Oil Crops Yearbook 2009*. Washington, D.C.: USDA Economic Research Services.
- ERS. 2011. Table 04; Soybean meal and Table 05; Soybean oil, In Oil Crops Yearbook 2009. Washington, D.C.: USDA Economic Research Service. Available at: http://usda.mannlib.cornell. cdu/MannUsda/viewDocumentInfo.do?documentID=1290. Accessed 29 March 2011.
- Huo, H., M. Wang, C. Bloyd, and V. Putsche. 2009. Life-cycle assessment of energy use and greenhouse gas emissions of soybean-derived biodiesel and renewable fuels. *Environ. Sci.* and Tech. 43(3): 750-756.

IndexMundi. 2012. Historical commodity prices. Available at:

- www.indexmundi.com/commodities/?commodity=soybeans&months=120. Accessed 10 May 2012.
- IPCC. 1996. IPCC Second Assessment Report: Climate Change 1995 (SAR). Working Group I: The Science of Climate Change. New York, N.Y.: Cambridge University Press. Available at: www.ipcc.ch/publications_and_data/publications_and_data_reports.shtml.
- IPCC. 2001. IPCC Third Assessment Report: Climate Change 2001 (TAR). Working Group I: The Scientific Basis. New York, N.Y.: Cambridge University Press. Available at www.ipcc.ch/ipccreports/tar/index.htm.
- IPCC. 2006. Chapter 11: N₂O emissions from managed soils, and CO₂ emissions from lime and urea application. In 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Vol.4: 11.1-11.54. Cambridge, U.K.: Cambridge University Press. Available at: www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_11_Ch11_N2O&CO2.pdf. Accessed 14 June 2011
- IPCC. 2007. Changes in atmospheric constituents and in radiative forcing. In *Climate Change 2007: The Physical Science Basis*, 130-234. S. Solomon, D. Qin, and M. Manning, ed. Cambridge, U.K.: Cambridge University Press.
- Landis, A. E., S. A. Miller, and T. L. Theis. 2007. Life cycle of the corn-soybean agroecosystem for biobased production. *Environ. Sci. Tech.* 41(4): 1457-1464.
- NASS. 2007. Agricultural chemical usage 2006 field crops summary. Ag Ch 1 (07) a. Washington, D.C.: USDA National Agricultural Statistics Service. Available at: http://usda.

- mannlib.cornell.edu/usda/nass/AgriChemUsFC//2000s/2007/ AgriChemUsFC-05-16-2007_revision.pdf. Accessed 14 June 2011.
- NASS. 2010. Quick Stats 2.0. Washington, D.C.: USDA National Agricultural Statistics Service. Available at: http://quickstats.nass.usda.gov/. Accessed 14 June 2011.
- Sheehan, J., V.Camobreco, J. Duffield, M. Graboski, and H. Shapouri. 1998. Life cycle inventory of biodiesel and petroleum diesel for use in an urban bus. NREL/SR-580-24089. Golden, Colo.: National Renewable Energy Laboratory.
- Thompson, J. C., and B. B. He. 2006. Characterization of crude glycerol from biodiesel production from multiple feedstocks. Applied Eng. in Agric. 22(2): 261-265.
- UNFCCC. 2006. Updated UNFCCC reporting guideline on annual inventories following incorporation of the provisions of decision 14/CP.11. FCCC/SBSTA/2006/9. Bonn, Germany: United Nations Framework Convention on Climate Change (UNFCCC). Available at: http://unfccc.int/resource/docs/2006/sbsta/eng/09.pdf. Accessed 14 June 2011.
- Van Gerpen, J., R. Pruszko, D. Celments, B. Shanks, and G. Knothe. 2006. Building a Successful Biodiesel Business: Technology Considerations, Developing the Business, Analytical Methodologies. 2nd ed. Dubuque, Iowa: Biodiesel Basies
- Woerfel, J. B. 1995. Chapter 6: Extraction. In Practical Handbook of Soybean Processing and Utilization, 65-92. D. R. Erickson, ed. Champaign, Ill.: AOCS Press and United Soybean Board.

2264 TRANSACTIONS OF THE ASABE

DRAFT Staff Summary: Mixed Feedstock for Renewable Diesel Method 2B Pathway Diamond Green Diesel

Plant Summary

Diamond Green Diesel (DGD) has submitted a Method 2B application for the production of mixed-feedstock renewable diesel (RD) at its St. Charles, Louisiana plant. The St. Charles plant, which is currently under construction, will be capable of producing 420,000 gallons of RD per day. DGD expects the plant to begin producing RD from the following seven feedstocks in early 2013: Midwestern soy oil, Midwestern corn oil, Midwestern used cooking oil (cooking required), Midwestern used cooking oil (no cooking), U.S. animal fat (higher energy), and U.S. animal fat (lower energy). All seven of DGD's pathways are modified versions of existing LCFS RD or biodiesel pathways. All are modeled as UOP Econofining Processes and utilize the default RD process energy consumption values found in CA-GREET 1.8b. Those process energy defaults are summarized in Table 1 below. All individual feedstocks present in the feedstock mixtures to be run at the St. Charles plant will be tracked by an inventory management system that is integrated into the plant's accounting system. The carbon intensity of all gallons of RD produced will be labeled with the CI of individual feedstocks, in keeping with the mixed-feedstock bio-and renewable diesel guidance published by ARB¹.

Table 1: Energy for Renewable Diesel Process from CA-GREET model

Feedstock	Process Energy Input (Btu/lb)	Electricity and Thermal Energy Shares (%Electric/%Thermal)
Soy Oil	1.051	
Corn Oil	1,851	61.4% / 38.6%
Used Cooking Oil (UCO)	2.475	01.470 / 30.070
Tallow	2,175	

Operating Conditions

Method 2 applications covering operating plants must base CI calculations on operational data covering two years, whenever possible. Because the DGD application covers a plant that is not yet operational, DGD will submit energy consumption data for the first two years of operation. Data submission will occur no less frequently than annually. If the data submitted indicates that any of DGD's actual production CIs are significantly higher than its approved LCFS pathway CIs, those CIs will be adjusted to better reflect actual operating conditions.

Page 1 of 4

¹ Air Resources Board, December 3, 2012. "Mixed-Feedstock Bio- and Renewable Diesel Guidance." http://www.arb.ca.gov/fuels/lcfs/2a2b/2a-2b-apps.htm.

As a condition of approval, DGD agrees to make all approved pathway CIs available via the LCFS Method 1 Lookup Tables to other RD producers whose production pathways are accurately described by the approved pathways developed in the DGD application.

Carbon Intensity of the Fuel Produced

Because all of DGD's pathway CIs are either higher than the corresponding reference pathways in the LCFS Method 1 Lookup table, or modified versions of LCFS renewable diesel pathways, its application falls under the Method 2B provisions of the LCFS. Method 2B applications are not subject to the substantiality requirements with which Method 2A applications must comply (a minimum improvement of five gCO₂e/MJ, and a minimum production volume of ten million gallons per year).

The proposed DGD pathway CIs are summarized in Table 2.

Page 2 of 4

Table 2: Proposed Lookup Table Entry

Fuel Pathway		Pathway	Carbon Intensity in gCO₂e/MJ (Including Indirect Effects)		
Fuel	Identifier	Description	Direct Emission	Land Use or Other Indirect Effect	Total
	RNWD 010	Conversion of Midwest soybean to renewable diesel (rail transport)	21.70	62	83.70
	RNWD 011	Conversion of Midwest soybean to soy oil to renewable diesel (ship transport)	21.48	62	83.48
	RNWD 012	Renewable diesel from Midwest corn oil produced from Dry DGS (rail transport)	6.00	0	6.00
	RNWD 013	Renewable diesel from Midwest corn oil produced from Dry DGS (ship transport)	5.56	0	5.56
Renewable Diesel	RNWD 016	Conversion of waste oils (Used Cooking Oil) from Midwest to renewable diesel where "cooking" is required (rail transport)	18.40	0	18.40
	RNWD 017	Conversion of waste oils (Used Cooking Oil) from Midwest to renewable diesel where "cooking" is required (ship transport)	18.18	0	18.18
	RNWD 018	Conversion of waste oils (Used Cooking Oil) from Midwest to renewable diesel where "cooking" is not required (rail transport)	13.85	0	13.85

Fuel	Pathway Identifier	Pathway Description		Intensity in gCo ding Indirect Eff	
	RNWD 019	Conversion of waste oils (Used Cooking Oil) from Midwest to renewable diesel where "cooking" is not required (ship transport)	13.63	0	13.63
	RNWD 020	Conversion of U.S. tallow to renewable diesel using higher energy use for rendering (rail transport)	40.34	0	40.34
	RNWD 021	Conversion of U.S. tallow to renewable diesel using higher energy use for rendering (ship transport)	40.12	0	40.12
	RNWD 022	Conversion of U.S. tallow to renewable diesel using lower energy use for rendering (rail transport)	19.91	0	19.91
	RNWD 023	Conversion of U.S. tallow to renewable diesel using lower energy use for rendering (ship transport)	19.70	0	19.70

Staff Analysis and Recommendation

Staff has reviewed DGD's application, and finds the following:

- Staff has replicated, using the CA-GREET spreadsheet, the carbon intensity values calculated by DGD; and
- Staff has confirmed that the energy consumption values used in the DGD application are the CA-GREET 1.8b defaults

On the basis of these findings, staff recommends that DGD's Method 2B pathways be approved for use in DGD's mixed-feedstock RD plant.

Biodiesel Profitability

Overview and Assumptions - Overview of the model, assumptions and data sources.

Economic Facility Model - The economic model that computes the monthly costs, revenue and profit (loss).

Tables:

Costs and Returns – Monthly Results per Gallon of Biodiesel
Costs and Returns – Monthly Results per Pound of Soybean Oil

To navigate among the pages in thi workbook, use the tabs at the botto of the page or the links in the text o this page.

Charts:

Input and Output Prices - Monthly biodiesel and glycerine prices -- 2007 to present.

Biodiesel Revenue - Monthly biodiesel and glycerine revenue - 2007 to present

Biodiesel Costs - Monthly cost to produce biodiesel per gallon (total and divided by category) - 2007 to present.

Biodiesel Revenue, Costs and Profits - Monthly costs and returns per gallon - 2007 to present.

Return on Equity - Monthly percent return on equity -- 2007 to present.

Breakeven Purchase Cost for Soybean oil and Sale Price for Biodiesel - Monthly prices facility can pay for soybean oil and receive

for biodiesel just to cover costs -- 2007 to present.

Biodiesel Revenue, Variable Costs and Profits - Monthly variable costs and returns per gallon - 2007 to present.

Ag Marketing Resource Center, Biodiesel Profitability

Authors: Don Hofstrand (dhof@iastate.edu)

IOWA STATE UNIVERS: Extension and Outreach

File updated: 11/06/2013 Next update: 12/15/2013

and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and mantal or family status. (Not all prohibited bases apply to all programs.) Many materials can be made available in alternative formats for ADA clients. To file a complaint of discrimination, write USDA, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and July 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa.

Overview and Assumptions of Biodiesel Profitability

The profitability of biodiesel prodution is extremely variable. Due to the volatile price nature of baseleland soybeanoil, its major feedstock, biodiesel profitability can change rapity from month to month. In addion, price variations of 6 co-product (glycering and its energy source (natural gas) add to the variativy of biodesel profits.

To see, the promoting of occasion production, an economic mode of a system one discrete pernings occasion. This is a so mallot gain, facility with construction costs simile to plants built in 2007. The costs and efficiencies are believed to be typical of lower biodisell plants. The prices of biodiesel, glycerine, soyber oil and natural gas are updated monthly to compate the current profrability of biodiesel production.

Monthly price variables

- 1) Biodiesel Price Weekly price F O B (Free on Board) the plant (conveted into monthly average prices) asreported in the National Weekly Ag Energy Round-up by the USDA Ag Marketing Service

 2) Soybean Oil Price Daily price converted no monthly average prices as reported by the USDAAg Market ResearchService, Jown Soybean Oil Price Monthly Average Reginal Posted Contract Price History repated by Methanex.

 3) Methanol Price Monthly Iowa natural ga price for indistrial users as reported by the Energy Information Administration (official energy statistics of the U.S. government).

Although these prices are representative of laws biodiesel plants, bey may not be representative of plants in other regions or states in the economic model the user can increase or decrease any 6the price series by a fixed amount to represent a special studion. An adjustment in a price series while reflected in the analysis tables and graphs.

Revenue costs and net returns (pr@tability) are shown monthly per gallon of biodiesel and per 00 pounds of soybean oil. Also, biodiesel and soy oil price breakwen levels are computed.

Major assumptions and characteristics of the biodiesel plant model

- 1) Turnkey biodesel production facility

- 1) Turney blooses production facility
 2) Facility but in 2007
 3) Nameplate capacity of 30 million gallons
 4) Facility construction cost (including working capital) of \$1.57 per gallon of namepate capacity
 5) Lender finance 50 percent of the project
 6) Equity financing of 50 percent of the project
 7) Plant operates at 100 percent of nameplate capacity
 8) Conversion factor of 7.55 pends of soybean oil per gallon of biodiesel

- 10) Agailon of biodiesel produces 9 pounds of glycerine
 10) Natural gas requirement of 7 cubic feet per gallon of biodiesel
 11) Typical input costs for an lowa soylban oil biodiesel facility.

The monthly profability of this hypothecal plant is computed by using the monthly markeprices for biodesel, soybean oil, methanol and natural gas. Each month the analysis is updated with the previous month's prices if any of these price data series do notify your situation, you can enter analysisment factor that will increase or decrese the coefficients in the price data series All other variables are held constant throughout the analysis.

Input coefficient adjustment. Allough we believe the coefficients in this model are a god representation of a soyban oil biodiesel plant, the user has the ability to change any of the inpd coefficients in the economic model to fit a special *uution. A change in an input coefficient will be reflected in the analysis tables and grap.

Ag Marketing Resource Center, Biodiesel Profitability

Authors Don Hofstrand & Ann Johanns

IOWA STATE UNIVERSITY Extension and Outreach

File updated 11/6/2013

The US Expertment AproAxis (ECA protests documentum in an impropers are tabled on the best offers code means organ perspending on April 4 posts offers and a restrict organization of the protection of the protec

Economic Model of a Biodiesel Production Facility Place the cursor over cells with red triangles to read comments.

Assumptions (inputs)

Facility (Construction		
Nameplate Capacity	30,000,000	gal./yr.	
Organizational Costs	\$200,000		
Process System	\$30,000,000		
Land, Site and Other	\$7,400,000		
Construction Related Costs	\$2,500,000		
Office and Administration	\$900,000		
Inventory & Working Capital	\$6,000,000		
Estimated Life	15	years	
Property Taxes	\$50,000	per yr.	

, <u>F</u>	inancing
Percent Debt	50 %
Length of Loan	10 years
Interest Rate	8.25 %

7.55 lbs./gal. biodiesel
1000/
100 % capacity
0.90 lbs./gal. biodiesel
0.71 lbs./gal. biodiesel
7 cub. ft./gal. biodiesel
0.6 Kwh./gal. biodiesel
2.0 gal./gal. biodiesel

Chemica	ıls .
Chemicals and Ingredients	5.70 ¢/gallon

	Prices	
Electricity	5.000 ¢/KwH	
Water	0.350 ¢/gallon	
Glycerin	3.000 ¢/pound	

Labor	& Management	
	Number	Salary
Operations	12	\$68,000
Maintenance	2	\$72,000
Laboratory	1	\$80,000
Material Handlers	10	\$30,000
Administration	3	\$90,000
Total	28	\$1,610,000

Other Dire	ct Costs		
Repairs & Maintenance	3.00	¢/gallon	
Transportation	10.00	¢/gallon	
Marketing and Procurement	4.00	¢/gallon	
Other	3.00	¢/gallon	

	Price Adjustments (+ or -)
Biodiesel	\$0.00 \$/gallon
Soybean Oil	0.00 ¢/gallon
Natural Gas	\$0.00 \$/1000 cubic feet

Output

Annual Production a	nd Resource Usage
Construction Cost	
Total	\$47,000,000
Equity	\$23,500,000
Debt	\$23,500,000
Per Gal. Nameplate Capacity	\$1.57
Per Gal. Operating Capacity	\$1.57
Per Lb. Soybean Oil Oper. Cap.	\$0.21
Depreciation	\$2,733,333 per year
Biodiesel Production	
Nameplate Capacity	30,000,000 gallons per year
Operating Capacity	30,000,000 gallons per year
Glycerine Production	27,000,000 pounds per year
Soybean Oil Usage	226,500,000 pounds per year
Natural Gas Usage	210,000 1,000 cubic feet/year
Electricity Usage	18,000,000 kilowatt hours/year
Electricity Cost	\$900,000 per year
Water Usage	60,000,000 gallons per year
Water Cost	\$210,000 per year
Number of Employees	28 employees
Labor & Management Cost	\$1,610,000 per year
Interest Cost	\$1,938,750 per year
Chemicals and Ingredients	\$1,710,000 per year
Repairs & Maintenance	\$200,000 per year
Transportation Cost	\$3,000,000 per year
Other Costs	\$200,000 per year

Produ	ction Costs	
Chemicals	Cost per Gallon	Cost per Pound
Chemicals and Ingredients	5.70 ¢/gal.	0.75¢/lb.
Total Chemical Cost	5.70 ¢/gal.	0.75¢/lb.
Other Direct Costs		
Repairs & Maintenance	3.00 ¢/gal.	0.40¢/lb.
Transportation	10.00 ¢/gal.	1.32¢/lb.
Water	0.70 ¢/gal.	0.09¢/lb.
Electricity	3.00 ¢/gal.	0.40¢/lb.
Other	3.00 ¢/gal.	0.40¢/lb.
Total Other Costs	19.70 ¢/gal.	2.61¢/lb:
Fixed Costs		
Depreciation	9.11 ¢/gal.	1.21¢/lb.
Interest	6.46 ¢/gal.	0.86¢/lb.
Labor & Management	5.37 ¢/gal.	0.71¢/lb.
Marketing & Procurement	4.00 ¢/gal.	0.53¢/lb.
Property Taxes, Insurance, etc.	1.17 ¢/gal.	0.15¢/lb.
Total Fixed Costs	26.11 ¢/gal.	3.46¢/lb.
Total Costs (less corn & natural g	as)	
Total Variable Costs	25.40 ¢/gal.	3.36¢/lb.
Total Variable & Fixed Costs	51.51 ¢/gal.	6.82¢/lb.

Monthly Costs and Returns per Gallon of Biodiesel Produced


		Prices			Revenue per Gallon			Cost per Gallon							Net Re		
Month		<u></u>	Natural					1	Natural		Other	Total			Biodiesel	Over	Over
and		Soybean Oil		Methanol	1000	Glycerine	Total	Soybean Oil	Gas	Methanol	Variable	Variable	Fixed	Total	Break-	Variable	All
Year	(gallon)		(000 cub. ft)	(pound)	Revenue	Revenue	Revenue	Cost	Cost	Cost	Costs	Costs	Costs	Cost	even	Costs	Costs
Apr-07	\$ 3.09	\$ 0.30	\$ 8.58	\$ 0.15	\$ 3.09	\$ 0.03	\$ 3.12	\$ 2.26	\$ 0.06	\$ 0.11	\$ 0.25	\$ 2.68	\$ 0.26	\$ 2.94	\$ 2.91	\$ 0.44	\$ 0.18
May-07	\$ 3.16	\$ 0.32	\$ 8.00	\$ 0.15	\$ 3.16	\$ 0.03	\$ 3.18	\$ 2.43	\$ 0.06	\$ 0.11	\$ 0.25	\$ 2.85	\$ 0.26	\$ 3.11	\$ 3.08	\$ 0.33	\$ 0.07
Jun-07	\$ 3.17	\$ 0.33	\$ 8.58	\$ 0.15	\$ 3.17	\$ 0.03	\$ 3.19	\$ 2.49	\$ 0.06	\$ 0.11	\$ 0.25	\$ 2.91	\$ 0.26	\$ 3.17	\$ 3.15	\$ 0.28	\$ 0.02
Jul-07	\$ 3.20	\$ 0.34	\$ 8.61	\$ 0.14	\$ 3.20	\$ 0.03	\$ 3.22	\$ 2.60	\$ 0.06	\$ 0.10	\$ 0.25	\$ 3.01	\$ 0.26	\$ 3.27	\$ 3.24	\$ 0.21	\$ (0.05)
ug-07	\$ 3.22	\$ 0.34	\$ 7.88	\$ 0.14	\$ 3.22	\$ 0.03	\$ 3.25	\$ 2.54	\$ 0.06	\$ 0.10	\$ 0.25	\$ 2.94	\$ 0.26	\$ 3.21	\$ 3.18	\$ 0.30	\$ 0.04
ep-07	\$ 3.29	\$ 0.37	\$ 7.48	\$ 0.14	\$ 3.29	\$ 0.03	\$ 3.32	\$ 2.77	\$ 0.05	\$ 0.10	\$ 0.25	\$ 3.18	\$ 0.26	\$ 3.44	\$ 3.41	\$ 0.14	\$ (0.12)
Oct-07	\$ 3.44	\$ 0.38	\$ 7.48	\$ 0.26	\$ 3.44	\$ 0.03	\$ 3.47	\$ 2.87	\$ 0.05	\$ 0.18	\$ 0.25	\$ 3.36	\$ 0.26	\$ 3.62	\$ 3.59	\$ 0.11	\$ (0.15)
lov-07	\$ 3.74	\$ 0.43	\$ 8.53	\$ 0.30	\$ 3.74	\$ 0.03	\$ 3.77	\$ 3.24	\$ 0.06	\$ 0.21	\$ 0.25	\$ 3.77	\$ 0.26	\$ 4.03	\$ 4.01	\$ (0.00)	\$ (0.26)
ec-07	\$ 3.92	\$ 0.44	\$ 8.86	\$ 0.38	\$ 3.92	\$ 0.03	\$ 3.95	\$ 3.35	\$ 0.06	\$ 0.27	\$ 0.25	\$ 3.94	\$ 0.26	\$ 4.20	\$ 4.17	\$ 0.01	\$ (0.25)
lan-08	\$ 4.28	\$ 0.49	\$ 8.74	\$ 0.38	\$ 4.28	\$ 0.03	\$ 4.30	\$ 3.71	\$ 0.06	\$ 0.27	\$ 0.25	\$ 4.29	\$ 0.26	\$ 4.55	\$ 4.53	\$ 0.01	\$ (0.25)
eb-08	\$ 4.68	\$ 0.56	\$ 9.99	\$ 0.32	\$ 4.68	\$ 0.03	\$ 4.70	\$ 4.26	\$ 0.07	\$ 0.22	\$ 0.25	\$ 4.81	\$ 0.26	\$ 5.07	\$ 5.05	\$ (0.11)	\$ (0.37)
Mar-08	\$ 5.16	\$ 0.57	\$ 10.06	\$ 0.29	\$ 5.16	\$ 0.03	\$ 5.19	\$ 4.32	\$ 0.07	\$ 0.20	\$ 0.25	\$ 4.85	\$ 0.26	\$ 5.11	\$ 5.08	\$ 0.34	\$ 0.08
Apr-08	\$ 4.98	\$ 0.56	\$ 10.71	\$ 0.24	\$ 4.98	\$ 0.03	\$ 5.01	\$ 4.21	\$ 0.07	\$ 0.17	\$ 0.25	\$ 4.71	\$ 0.26	\$ 4.97	\$ 4.94	\$ 0.30	\$ 0.04
lay-08	\$ 5.26	\$ 0.58	\$ 10.89	\$ 0.23	\$ 5.26	\$ 0.03	\$ 5.28	\$ 4.38	\$ 0.08	\$ 0.16	\$ 0.25	\$ 4.87	\$ 0.26	\$ 5.13	\$ 5.10	\$ 0.42	\$ 0.15
un-08	\$ 5.51	\$ 0.62	\$ 11.83	\$ 0.24	\$ 5.51	\$ 0.03	\$ 5.54	\$ 4.65	\$ 0.08	\$ 0.17	\$ 0.25	\$ 5.16	\$ 0.26	\$ 5.42	\$ 5.39	\$ 0.38	\$ 0.12
Jul-08	\$ 5.47	\$ 0.60	\$ 12.42	\$ 0.24	\$ 5.47	\$ 0.03	\$ 5.49	\$ 4.54	\$ 0.09	\$ 0.17	\$ 0.25	\$ 5.05	\$ 0.26	\$ 5.31	\$ 5.28	\$ 0.45	\$ 0.19
ug-08	\$ 4.88	\$ 0.51	\$ 11.83	\$ 0.24	\$ 4.88	\$ 0.03	\$ 4.90	\$ 3.82	\$ 0.08	\$ 0.17	\$ 0.25	\$ 4.33	\$ 0.26	\$ 4.59	\$ 4.56	\$ 0.57	\$ 0.31
ep-08	\$ 4.43	\$ 0.46	\$ 9.30	\$ 0.24	\$ 4.43	\$ 0.03	\$ 4.46		\$ 0.07	\$ 0.17	\$ 0.25	\$ 3.93	\$ 0.26	\$ 4.19	\$ 4.16	\$ 0.53	\$ 0.27
ct-08	\$ 3.65	\$ 0.35	\$ 7.30	\$ 0.23	\$ 3.65	\$ 0.03	\$ 3.67		\$ 0.05	\$ 0.16	\$ 0.25	\$ 3.10	\$ 0.26	\$ 3.36	\$ 3.33	\$ 0.57	\$ 0.31
ov-08	\$ 3.19	\$ 0.32	\$ 7.11	\$ 0.21	\$ 3.19	\$ 0.03	\$ 3.22		\$ 0.05	\$ 0.15	\$ 0.25	\$ 2.86	\$ 0.26	\$ 3.12	\$ 3.09	\$ 0.36	\$ 0.10
ec-08	\$ 2.84	\$ 0.29	\$ 7.92	\$ 0.15	\$ 2.84	\$ 0.03	\$ 2.87	LIGOTON CO.	\$ 0.06	\$ 0.11	\$ 0.25	\$ 2.58	\$ 0.26	\$ 2.84	\$ 2.82	\$ 0.29	\$ 0.03
an-09	\$ 3.09	\$ 0.32	\$8.22	\$ 0.11	\$ 3.09	\$ 0.03	\$ 3.12		\$ 0.06	\$ 0.08	\$ 0.25	\$ 2.78	\$ 0.26	\$ 3 04	\$ 3.02	\$ 0.33	\$ 0.07
eb-09	\$ 2.82	\$ 0.29	\$ 7.84	\$ 0.11	\$ 2.82	\$ 0.03	\$ 2.85	USP30-00-10 R	\$ 0.05	\$ 0.08	\$ 0.25	\$ 2.56	\$ 0.26	\$ 2.82	\$ 2.79	\$ 0.29	\$ 0.03
1ar-09	\$ 2.68	\$ 0.28	\$ 7.28	\$ 0.10	\$ 2.68	\$ 0.03	\$ 2.71	1 15 15 15 15 15 15	\$ 0.05	\$ 0.07	\$ 0.25	\$ 2.49	\$ 0.26	\$ 2.76	\$ 2.73	\$ 0.21	\$ (0.05)
pr-09	\$ 2.94	\$ 0.33	\$ 5.60	\$ 0.09	\$ 2.94	\$ 0.03	\$ 2.97		\$ 0.04	\$ 0.06	\$ 0.25	\$ 2.83	\$ 0.26	\$ 3.09	\$ 3.06	\$ 0.14	\$ (0.12)
lay-09	\$ 3.10	\$ 0.36	\$ 4.84	\$ 0.09	\$ 3.10	\$ 0.03	\$ 3.13		\$ 0.03	\$ 0.06	\$ 0.25	\$ 3.09	\$ 0.26	\$ 3.35	\$ 3.32	\$ 0.04	\$ (0.22)
un-09	\$ 3.13	\$ 0.35	\$ 4.57	\$ 0.09	\$ 3.13	\$ 0.03	\$ 3.16	1.1000000000000000000000000000000000000	\$ 0.03	\$ 0.06	\$ 0.25	\$ 3.03	\$ 0.26	\$ 3.29	\$ 3.26	\$ 0.13	\$ (0.13)
Jul-09	\$ 2.86	\$ 0.31	\$ 4.55	\$ 0.10	\$ 2.86	\$ 0.03	\$ 2.89		\$ 0.03	\$ 0.00	\$ 0.25	\$ 2.73	\$ 0.26	\$ 2.99	\$ 2.96	\$ 0.16	\$ (0.10)
ug-09	\$ 3.11	\$ 0.34	\$ 4.73	\$ 0.11	\$ 3.11	\$ 0.03	\$ 3.13		\$ 0.03	\$ 0.08	\$ 0.25	\$ 2.90	\$ 0.26	\$ 3.16	\$ 3.13	\$ 0.24	\$ (0.02)
ep-09	\$ 3.02	\$ 0.31	\$ 4.68	\$ 0.13	\$ 3.02	\$ 0.03	\$ 3.04		\$ 0.03	\$ 0.09	\$ 0.25	\$ 2.72	\$ 0.26	\$ 2.99	\$ 2.96	\$ 0.32	\$ 0.06
ct-09	\$ 3.12	\$ 0.34	\$ 4.84	\$ 0.14	\$ 3.12	\$ 0.03	\$ 3.15		\$ 0.03	\$ 0.10	\$ 0.25	\$ 2.93	\$ 0.26	\$ 3.19	\$ 3.17	\$ 0.21	\$ (0.05)
ov-09	\$ 3.37	\$ 0.37	\$ 6.16	\$ 0.15	\$ 3.37	\$ 0.03	\$ 3.40		\$ 0.03	\$ 0.10	\$ 0.25	\$ 3.16	\$ 0.26	\$ 3.42	\$ 3.39	\$ 0.24	\$ (0.03)
ec-09	\$ 3.36	\$ 0.37	\$ 6.79	\$ 0.17	\$ 3.36	\$ 0.03	\$ 3.39		\$ 0.04	\$ 0.12	\$ 0.25	\$ 3.21	\$ 0.26	\$ 3.48	\$ 3.45	\$ 0.17	
an-10	\$ 3.31	\$ 0.36	\$ 6.61	\$ 0.17	\$ 3.31	\$ 0.03	\$ 3.34		\$ 0.05	\$ 0.12	\$ 0.25	\$ 3.10	\$ 0.26	\$ 3.36	\$ 3.43	\$ 0.24	\$ (0.09)
eb-10	\$ 3.29	\$ 0.35	\$ 7.08	\$ 0.17	\$ 3.29	\$ 0.03	\$ 3.32	\$1000 (CONT.)	\$ 0.05	\$ 0.12	\$ 0.25	\$ 3.10	\$ 0.26	\$ 3.35	\$ 3.33		
ar-10	\$ 3.31	\$ 0.36	\$ 7.06	\$ 0.17	\$ 3.31					2.5 2.5 d 2.			(T) () (T) (T) (T)	12/00/2012	100000000000000000000000000000000000000	\$ 0.23	\$ (0.03)
pr-10	\$ 3.32	\$ 0.36	\$ 6.21	\$ 0.17	\$ 3.31	\$ 0.03	\$ 3.34		\$ 0.05	\$ 0.12	\$ 0.25	\$ 3.18	\$ 0.26	\$ 3.44	\$ 3.41	\$ 0.17	\$ (0.10)
	\$ 3.32	\$ 0.35	150507000			\$ 0.03	\$ 3.35		\$ 0.04	\$ 0.12	\$ 0.25	\$ 3.16	\$ 0.26	\$ 3.42	\$ 3.39	\$ 0.19	\$ (0.07)
ay-10_ un-10	\$ 3.23	\$ 0.35	\$ 5.45	\$ 0.15	\$ 3.23	\$ 0.03	\$ 3.26		\$ 0.04	\$ 0.11	\$ 0.25	\$ 3.05	\$ 0.26	\$ 3.31	\$ 3.29	\$ 0.21	\$ (0.05)
	\$ 3.17		\$ 5 61	\$ 0.16	\$ 3.17	\$ 0.03	\$ 3.20		\$ 0.04	\$ 0.11	\$ 0.25	\$ 3.03	\$ 0.26	\$ 3.29	\$ 3.26	\$ 0.17	\$ (0.09)
Jul-10_	10.50	\$ 0.35	\$ 5.69	\$ 0.16	\$ 3.18	\$ 0.03	\$ 3.21		\$ 0.04	\$ 0.11	\$ 0.25	\$ 3.07	\$ 0.26	\$ 3.33	\$ 3.30	\$ 0.14	\$ (0.12)
ug-10_	\$ 3.30	\$ 0.36	\$ 5.76	\$ 0.16	\$ 3.30	\$ 0.03	\$ 3.33		\$ 0.04	\$ 0.11	\$ 0.25	\$ 3.16	\$ 0.26	\$ 3.42	\$ 3.39	\$ 0.17	\$ (0.09)
ep-10	\$ 3.35	\$ 0.39	\$ 5.59	\$ 0.16	\$ 3.35	\$ 0.03	\$ 3.38	\$ 2.95	\$ 0.04	\$ 0.12	\$ 0.25	\$ 3.35	\$ 0.26	\$ 3.62	\$ 3.59	\$ 0.02	\$ (0.24)

2634


Monthly Biodiesel Costs and Returns per Pound of Soybean Oil Processed

20 23		Pr	rices		Revenue per Pound				To	tal Cost/LI	o.	Net Ret	Return	
Month	Natural							Soybean Oil				Over Over		on
and	Biodiesel	Soybean O	il Gas	Methanol	Biodiesel	Glycerine	Total	Break-	Variable	Fixed	Total	Variable	All	Equity
Year	(per gal.)	(per lbs.)	(000 cub. ft.)	(per lbs.)	Revenue	Revenue	Revenue	even	Costs	Costs	Costs	Costs	Costs	Annual
Apr-07	\$ 3.09	\$ 0.30	\$ 8.58	\$ 0.15	\$ 0.41	\$ 0.00	\$ 0.41	\$ 0.32	\$ 0.35	\$ 0.03	\$ 0.39	\$ 0.06	\$ 0.02	23%
May-07	\$ 3.16	\$ 0.32	\$ 8.00	\$ 0.15	\$ 0.42	\$ 0.00	\$ 0.42	\$ 0.33	\$ 0.38	\$ 0.03	\$ 0.41	\$ 0.04	\$ 0.01	9%
Jun-07	\$ 3.17	\$ 0.33	\$ 8.58	\$ 0.15	\$ 0.42	\$ 0.00	\$ 0.42	\$ 0.33	\$ 0.39	\$ 0.03	\$ 0.42	\$ 0.04	\$ 0.00	2%
Jul-07	\$ 3.20	\$ 0.34	\$ 8.61	\$ 0.14	\$ 0.42	\$ 0.00	\$ 0.43	\$ 0.34	\$ 0.40	\$ 0.03	\$ 0.43	\$ 0.03	\$ (0.01)	-6%
Aug-07	\$ 3.22	\$ 0.34	\$ 7.88	\$ 0.14	\$ 0.43	\$ 0.00	\$ 0.43	\$ 0.34	\$ 0.39	\$ 0.03	\$ 0.42	\$ 0.04	\$ 0.01	5%
Sep-07	\$ 3.29	\$ 0.37	\$ 7.48	\$ 0.14	\$ 0.44	\$ 0.00	\$ 0.44	\$ 0.35	\$ 0.42	\$ 0.03	\$ 0.46	\$ 0.02	\$ (0.02)	-16%
Oct-07	\$ 3.44	\$ 0.38	\$ 7.48	\$ 0.26	\$ 0.46	\$ 0.00	\$ 0.46	\$ 0.36	\$ 0.44	\$ 0.03	\$ 0.48	\$ 0.01	\$ (0.02)	-19%
Nov-07	\$ 3.74	\$ 0.43	\$ 8.53	\$ 0.30	\$ 0.50	\$ 0.00	\$ 0.50	\$ 0.39	\$ 0.50	\$ 0.03	\$ 0.53	\$ (0.00)	\$ (0.03)	-34%
Dec-07	\$ 3.92	\$ 0.44	\$ 8.86	\$ 0.38	\$ 0.52	\$ 0.00	\$ 0.52	\$ 0.41	\$ 0.52	\$ 0.03	\$ 0.56	\$ 0.00	\$ (0.03)	-32%
Jan-08	\$ 4.28	\$ 0.49	\$ 8.74	\$ 0.38	\$ 0.57	\$ 0.00	\$ 0.57	\$ 0.46	\$ 0.57	\$ 0.03	\$ 0.60	\$ 0.00	\$ (0.03)	-32%
Feb-08	\$ 4.68	\$ 0.56	\$ 9.99	\$ 0.32	\$ 0.62	\$ 0.00	\$ 0.62	\$ 0.52	\$ 0.64	\$ 0.03	\$ 0.67	\$ (0.01)	\$ (0.05)	-47%
Mar-08	\$ 5.16	\$ 0.57	\$ 10.06	\$ 0.29	\$ 0.68	\$ 0.00	\$ 0.69	\$ 0.58	\$ 0.64	\$ 0.03	\$ 0.68	\$ 0.05	\$ 0.01	10%
Apr-08	\$ 4.98	\$ 0.56	\$ 10.71	\$ 0.24	\$ 0.66	\$ 0.00	\$ 0.66	\$ 0.56	\$ 0.62	\$ 0.03	\$ 0.66	\$ 0.04	\$ 0.00	5%
May-08	\$ 5.26	\$ 0.58	\$ 10.89	\$ 0.23	\$ 0.70	\$ 0.00	\$ 0.70	\$ 0.60	\$ 0.64	\$ 0.03	\$ 0.68	\$ 0.06	\$ 0.02	20%
Jun-08	\$ 5.51	\$ 0.62	\$ 11.83	\$ 0.24	\$ 0.73	\$ 0.00	\$ 0.73	\$ 0.63	\$ 0.68	\$ 0.03	\$ 0.72	\$ 0.05	\$ 0.02	15%
Jul-08	\$ 5.47	\$ 0.60	\$ 12.42	\$ 0.24	\$ 0.72	\$ 0.00	\$ 0.73	\$ 0 63	\$ 0.67	\$ 0.03	\$ 0.70	\$ 0.06	\$ 0.02	24%
Aug-08	\$ 4.88	\$ 0.51	\$ 11.83	\$ 0.24	\$ 0.65	\$ 0.00	\$ 0.65	\$ 0.55	\$ 0.57	\$ 0.03	\$ 0.61	\$ 0.08	\$ 0.04	40%
Sep-08	\$ 4.43	\$ 0.46	\$ 9.30	\$ 0.24	\$ 0.59	\$ 0.00	\$ 0.59	\$ 0.49	\$ 0.52	\$ 0.03	\$ 0.55	\$ 0.07	\$ 0.04	35%
Oct-08	\$ 3.65	\$ 0.35	\$ 7.30	\$ 0.23	\$ 0.48	\$ 0.00	\$ 0.49	\$ 0.39	\$ 0.41	\$ 0.03	\$ 0.44	\$ 0.08	\$ 0.04	40%
Nov-08	\$ 3.19	\$ 0.32	\$ 7.11	\$ 0.21	\$ 0.42	\$ 0.00	\$ 0.43	\$ 0.33	\$ 0.38	\$ 0.03	\$ 0.41	\$ 0.05	\$ 0.01	13%
Dec-08	\$ 2.84	\$ 0.29	\$ 7.92	\$ 0.15	\$ 0.38	\$ 0.00	\$ 0.38	\$ 0.29	\$ 0.34	\$ 0.03	\$ 0.38	\$ 0.04	\$ 0.00	3%
Jan-09	\$ 3.09	\$ 0.32	\$ 8.22	\$ 0.11	\$ 0.41	\$ 0.00	\$ 0.41	\$ 0.33	\$ 0.37	\$ 0.03	\$ 0.40	\$ 0.04	\$ 0.01	9%
Feb-09	\$ 2.82	\$ 0.29	\$ 7.84	\$ 0.11	\$ 0.37	\$ 0.00	\$ 0.38	\$ 0.29	\$ 0.34	\$ 0.03	\$ 0.37	\$ 0.04	\$ 0.00	4%
Mar-09	\$ 2.68	\$ 0.28	\$ 7.28	\$ 0.10	\$ 0.36	\$ 0.00	\$ 0.36	\$ 0.27	\$ 0.33	\$ 0.03	\$ 0.36	\$ 0.03	\$ (0.01)	-6%
Apr-09	\$ 2.94	\$ 0.33	\$ 5.60	\$ 0.09	\$ 0.39	\$ 0.00	\$ 0.39	\$ 0.31	\$ 0.37	\$ 0.03	5 0 41	\$ 0.02	\$ (0.07)	-16%


2634




Percent Return on Equity (annualized basis) (2007 to present)



Total Revenue per Gallon (biodiesel, glycerine and total, 2007 to present)







Soybean Oil Breakeven and Market Price (Soybean Oil Breakeven = total revenue less all costs except soybean oil cost) (2007 to present)




Cost of Biodiesel Production (\$/gallon) (2007 to present)



Biodiesel Revenue, Variable Cost, and Return Over Variable Cost (\$ per gallon) (2007 - Present)



Biodiesel Revenue , Costs , and Profit (\$ per gallon) (2007 - Present)





Friday, March 26, 2010

Book 2 of 2 Books Pages 14669-15320

Part II

Environmental Protection Agency

40 CFR Part 80
Regulation of Fuels and Fuel Additives:
Changes to Renewable Fuel Standard
Program; Final Rule

Attachment 2010-3851. pdf Public Document

1	スプ
7	, ,
25.	Š
16	7
1	۲)
(DO
	8
	2
	36

iodiesel (billion (allons)	Feedstock Usage (22M data)	2013 feedstock % of industry production (22M data)	LCA Source	Feedstock % GHG reduction (LCA category)	Diesel Baseline emissions gCO2e/MJ	Biodiesel Carbon Intensity gCO2e/MJ	Biodiesel lb CO2e/125,000 BTU	GHG reduction in lbs CO2e/gallon of biodiesel	GHG reduction in billion lbs CO2e	GHG reduction in million tons CO2e	GHG reduction in million Metric tons CO2e
1.7 sc	•	48%	USEPA 2010	57.0% soy	89.7	38.6	11.2	14.1	23.9	11.9	10.8
	orn oil	10%	USEPA 2010	86.0% waste grease	89.7	12.6	3.7	21.2	36.0	18.0	16.4
	sed cooking oil	8%	USEPA 2010	86.0% waste grease	89.7	12.6	3.7				
	ther recycled grease	3%	USEPA 2010	86.0% waste grease	89.7	12.6	3.7				-
	nimal fat	11%	USEPA 2010	86.0% waste grease	89.7	12.6	3.7				
	enewable diesel	15%	USEPA 2010	86.0% waste grease	89.7	12.6	3.7				
1.7 ca	anola	5%	USEPA 2010	50.5% canola	89.7	44.4	12.9	12.5	21.2	10.6	9.
1.7		100%	USEPA 2010 weighted by 2013 industry production USDA/Idaho 2012	70.3% 2013 industry	89.7 89.7	26.6 21.2	7.7 6.2	17.3		14.7	
1.7 sc	ру	48%	USDA/Idaho 2012	77.9% soy	96	21.2	6.2	20.6	35.0	17.5	15.
1.7 cc	orn oil	10%	CARB 2011	95.8% corn oil	96	4.0	1.2	25.3	43.0	21.5	-
1.7 us	sed cooking oil	8%	USEPA 2010	86.9% waste grease	96	12.6	3.7	22.9		19.5	
1.7 ot	ther recycled grease	3%	USEPA 2010	86.9% waste grease	96	12.6	3.7				
1.7 ar	nimal fat	11%	USEPA 2010	86.9% waste grease	96	12.6	3.7				
10000	enewable diesel	15%	CARB 2011	79.5% tallow RD	96	19.7	5.7	21.0	35.7	17.8	16.
1.7 ca	anola	5%	USEPA 2010	53.7% canola	96	44.4	12.9	14.2	24.1	12.1	10.
1.7			best available data weighted by 2013 industry production	80.7% 2013 industry	96	18.5	5.4	21.3	36.2	18.1	16.

green = inputs can be changed

inputs from reference material calculated by table

	nput	Unit	2011	2012	2013
Feedstock Inp					
	Canola Oil	%	12%	11%	
	Distillers Corn Oil	%	4%	8%	The second
Feedstock	Soybean Oil	%	55%	52%	48%
Breakdown	Animal Fats	%	16%	13%	11%
	Yellow Grease	%	9%	9%	11%
	Renewable Diesel	%	4%	8%	15%
	Canola Oil	cents/pound	61.0	61.2	56.2
	Distillers Corn Oil	cents/pound	43.6	38.0	34.6
Cost of	Soybean Oil	cents/pound	54.1	51.7	47.9
Feedstocks	Animal Fats	cents/pound	47.1	42.5	38.6
	Yellow Grease	cents/pound	43.1	37.3	34.5
	Renewable Diesel	cents/pound	45.1	39.9	35.6
Required amount of	f feedstock	lb/gallon	7.5	7.5	7.5
Average Price of F	eedstock	\$/lb	\$ 0.52	\$ 0.49	\$ 0.43
Cost of Feedstock		\$/Gallon	\$ 3.90	\$ 3.66	\$ 3.20
OOST OF F CCUSTOCK	per Gallott	Ψ/Οαιίοπ	ψ 3.90	Ψ 3.00	Ψ 3.20
Carban Cantan	4 1				
Carbon Conter	10 - 00 - 00 - 00 - 00 - 00 - 00 - 00 -	Tax.			
	Canola Oil	%	53%	53%	54%
	Distillers Corn Oil	%	96%	96%	96%
Carbon Reduction	Soybean Oil	%	78%	78%	78%
by Percent	Animal Fats	%	87%	87%	87%
	Yellow Grease	%	87%	87%	87%
	Renewable Diesel	%	79%	79%	80%
Discal Carbon Inter	a a i b	I=000=041	04.40	05.04	00.00
Diesel Carbon Inter	isity	gCO2e/MJ	94.43	95.21	96.00
	Canola Oil	lb CO2e / gal	13.8	14.0	14.2
	Distillers Corn Oil	lb CO2e / gal	24.9	25.1	25.3
Carbon Reduction		lb CO2e / gal	20.2	20.4	20.6
in CO2e	Animal Fats	lb CO2e / gal	22.5	22.7	23.0
111 0020	Yellow Grease	lb CO2e / gal	22.5	22.7	23.0
1	Renewable Diesel	lb CO2e / gal	20.6	20.8	21.0
	Treflewable Diesel	In OOZe / gai	20.0	20.0	21.0
Average Carbon Re	eduction for Biodiesel	lb/gallon	20.2	20.8	21.3
Output					
Fixed Cost		\$/Gallon	\$0.26	\$0.26	\$0.26
Other Variable Cos		\$/Gallon	\$0.44	\$0.43	\$0.46
Value of Co-Produc		\$/Gallon	-\$0.03	-\$0.03	-\$0.03
Cost of Production	n	\$/Gallon	\$4.57	\$4.32	\$3.89
		Ta. 0. 11			
Diesel Price		\$/Gallon	\$3.05	\$3.11	\$3.01
Biodiesel cost differ	ential	\$/Gallon	\$1.52	\$1.21	\$0.88
Cost of Carbon		\$/Tonne	\$166.14	\$127.69	\$90.92

Conversion Units					
	grams/pound				
118,296	Btu/gallon of diesel				
0.001055	Btu/MJ				
2,205	Pounds/metric ton				

Feedstock Inputs

Sources: Feedstock breakdown values based on EIA data. The required amount of feedstock is derived from the EIA's breakdown of feedstock conversion

Carbon Content Inputs

Sources: Based on EPA estimates, supplemented by an USDA study and CARB.

Output

Sources: The estimated non-feedstock variable costs and fixed costs come from the lowa State production cost model of a standard plant in lowa producing biodiesel from soybean oil

Sources: The diesel prices are simple averages of spot price data from EIA: Los Angeles, CA Ultra-Low Sulfur CARB Diesel Spot Price (Dollars per Gallon)

PRELIMINARY DRAFT DISTRIBUTED FOR PUBLIC COMMENT November 4, 2011

Production of Biodiesel from Corn Oil Extraction at at Dry Mill Corn Ethanol Plants

Summary

ARB staff has developed a California corn oil biodiesel (BD) pathway in which the feedstock is produced in Midwestern corn ethanol plants and shipped to California for fuel production. The resulting pathway CI is 4.00 grams of CO₂-equivalent greenhouse gas emissions per mega joule of biodiesel produced (gCO₂e/MJ). Although the feedstock transport, biodiesel production, and finished fuel transport portions of this pathway are identical to those found in ARB's soybean-to-biodiesel pathway, the feedstock production portion has no precedent in any other pathway. Calculation of the CI for that step requires that the energy consumption and greenhouse gas (GHG) generation associated with the production of corn oil be appropriately allocated between corn ethanol and corn oil.

In order to begin co-producing corn oil, standard dry mill corn ethanol plants need only be retrofitted with a centrifuge-based extraction system. This system extracts corn oil from the distillers' grains that emerge from the fermentation and distillation processes. As such, it has no direct impacts on the production of corn ethanol. It does, however, reduce the volume and lipid content of the distillers' grains with solubles (DGS) the plant produces.

The corn oil that is extracted from the DGS stream is an unrefined product that has two primary uses: a livestock feed additive and a biodiesel feedstock. This document summarizes a California Low Carbon Fuel Standard (LCFS) pathway in which corn oil produced at dry mill corn ethanol plants is used to produce biodiesel. This pathway does not apply to production processes in which the extracted corn oil is used for purposes other than the production of biodiesel for use as a transportation fuel. In addition, it is specific to ethanol production environments in which all DGS is fully dried. Dry DGS has a moisture content of around ten percent.

ARB staff's estimate of the carbon intensity (CI) of corn oil biodiesel is based on information made available by Greenshift Inc.—a company that has commercialized a corn oil extraction process. Although Greenshift's is not the only available extraction process, more information is publicly available on its process than is available on alternative systems.

Under the Greenshift process, corn oil is removed from the DGS process stream through a combination of washing and centrifuging. The extracted corn oil is shipped to a biodiesel production plant where it is converted to fatty acid methyl

Page 1 of 2

¹ ARB (2009). Detailed California-Modified GREET Pathway for Conversion of Midwest Soybeans to Biodiesel (Fatty Acid Methyl Esters-FAME); version 3.0: http://www.arb.ca.gov/fuels/lcfs/121409lcfs_soybean.pdf

PRELIMINARY DRAFT DISTRIBUTED FOR PUBLIC COMMENT November 4, 2011

ester (FAME) biodiesel using a transesterification process—the same process that is used to produce biodiesel from soy oil and other oil- and fat-based feedstocks.

The equipment used to extract corn oil at corn ethanol plants consumes both thermal and electrical energy. This additional energy consumptions is more than offset, however, by energy savings realized during the DGS drying process. Energy is saved because the removal of corn oil both reduces the mass of the DGS entering the dryers, and improves the efficiency with which that DGS transfers heat. Based on information from Greenshift, ARB staff estimates that the production of corn oil at an ethanol plant reduces the net energy consumption of that plant by about nine percent. These savings would be realized only when all DGS is fully dried.

At dry mill plants that produce both ethanol and corn oil, the primary product is ethanol. Staff has no reason to believe that corn oil will ever replace ethanol as the primary product at such plants. Since corn oil production is incremental and secondary to ethanol production, staff has concluded that no portion of the GHG gas emissions associated with the production of ethanol should be allocated to corn oil biodiesel. Because corn oil extraction equipment can be installed in existing corn ethanol plants, ARB staff believes that the carbon intensity of corn oil should be calculated as a marginal, or incremental, carbon intensity, consisting only the additional energy requirements and savings that occur as a result of operating corn oil extraction equipment.

Staff is confident that corn oil biodiesel produced according to the pathway summarized above would have a carbon intensity of 4.00 gCO₂e/MJ. For that reason, staff recommends that the Executive Officer approve that pathway.

Page 2 of 2



Monthly Biodiesel Production Report

December 2013

With Data for October 2013

Independent Statistics & Analysis www.eia.gov

U.S. Department of Energy Washington, DC 20585



Biodiesel Highlights and Background, Data for October 2013

Highlights

Production – U.S. production of biodiesel was 132 million gallons in October 2013. Biodiesel production during October 2013 was about 5 million gallons higher than production in September 2013. Biodiesel production from the Midwest region (Petroleum Administration for Defense District 2) was 67% of the U.S. total. Production came from 112 biodiesel plants with capacity of 2.2 billion gallons per year.

Sales – Producer sales of biodiesel during October 2013 included 92 million gallons sold as B100 (100% biodiesel) and an additional 41 million gallons of B100 sold in biodiesel blends with diesel fuel derived from petroleum.

Feedstocks – There were a total of 1,009 million pounds of feedstocks used to produce biodiesel in October 2013. Soybean oil remained the largest biodiesel feedstock during October 2013 with 551 million pounds consumed.

Background

The Monthly Biodiesel Production Report provides data on operations of the U.S. biodiesel industry as part of EIA's response to section 1508 of the Energy Policy Act of 2005 which directed EIA to publish information on renewable fuels including biodiesel. Data are provided for the U.S. and in selected cases by state and region.

The source of data is <u>Form EIA-22M Monthly Biodiesel Production Survey</u>, used to collect the following information from registered producers of biodiesel.

- plant location, operating status, and annual production capacity
- production of 100% biodiesel (B100)
- · biodiesel coproduct production
- stocks
- · feedstock, alcohol input, and catalysts used in biodiesel production
- · sales of B100 and blended biodiesel from producing plants
- · sales of biodiesel to end-users

Form EIA-22M provides data necessary to monitor growth of the biodiesel industry in order to allow Congress to assess whether objectives of Section 503 of the Energy Policy Act of 1992 and Section 1508 of the Energy Policy Act of 2005 are being achieved.

Table 1. U.S. Biodiesel Production Capacity and Production (million gallons)

Period	Annual Production Capacity	Monthly B100 Productio
2011		
January	2,114	35
February	2,104	40
March	2,081	60
April	2,101	71
May	2,064	77
June	2,069	81
July	1,958	92
August	2,008	95
September	2,087	96
October	2,119	105
November	2,098	105
December	2,090	109
Total		967
2012		
January	2,203	74
February	2,203	79
March	2,188	95
April	2,158	94
May	2,151	102
June	2,138	93
July	2,134	89
August	2,135	91
September	2,165	82
October	2,162	75
November	2,091	57
December	2,054	59
Total		991
2013		
anuary	2,086	66
February	2,090	68
March	2,160	98
April	2,162	106
Мау	2,165	111
une	2,148	113
uly	2,144	128
August	2,141	128
September	2,153	127
October	2,190	132
0 Month Total	•	1,076
012 10 Month Total		874
011 10 Month Total		753
- = Not Applicable		

^{-- =} Not Applicable

Totals may not equal the sum of components due to independent rounding. B100 is the industry designation for pure biodiesel; a biodiesel blend contains both pure biodiesel and petroleum diesel fuel.

Source: U.S. Energy Information Administration, Form EIA-22M "Monthly Biodiesel Production Survey"

U.S. Energy Information Administration | Monthly Biodiesel Production Report

R = Revised

Table 2. U.S. Biodiesel Production, Sales, and Stocks (million gallons)

Period	B100 Production	Sales of B100	Sales of B100 Included in Biodiesel Blends	Ending Stocks of B100	B100 Stock Change
2011					
January	35	22	9	17	4
February	40	27	13	17	1
March	60	41	17	19	2
April	71	47	22	21	2
May	77	50	27	23	2
June	81	62	24	19	(4)
July	92	66	22	23	4
August	95	64	30	25	2
September	96	68	37	21	(4)
October	105	69	36	25	4
November	105	66	39	26	1
December	109	59	62	14	(12)
Total	967	641	339	-	1
2012		H. 184401.740 (1950)	Salas was in contract and a salas and		
January	74	45	16	26	12
February	79	57	15	33	7
March	95	75	17	36	2
April	94	68	24	40	5
May	102	77	33	37	(3)
June	93	80	19	31	(6)
July	89	72	18	29	(1)
August	91	73	18	30	1
September	82	62	17	32	2
October	75	62	12	33	1
November	57	53	12	26	(8)
December	59	44	10	31	5
Total	991	767	212	••	16
2013					
January	66	52	15	31	(1)
February	68	44	23	33	2
March	98	68	29	35	2
April	106	79	31	34	(1)
May	111	77	40	28	(6)
June	113	77	36	28	1
July	128	87	40	30	2
August	128	91	40	28	(2)
September	127	92	37	27	(1)
October	132	92	41	26	(1)
10 Month Total	1,076	758	331		(7)
2012 10 Month Total	874	670	189		19
2011 10 Month Total	753	516	238	••	13

^{-- =} Not Applicable

R = Revised

⁽s) = Value is less than 0.5 of the table metric, but value is included in any associated total.

Totals may not equal the sum of components due to independent rounding.

B100 is the industry designation for pure biodiesel; a biodiesel blend contains both pure biodiesel and petroleum diesel fuel.

Source: U.S. Energy Information Administration, Form EIA-22M "Monthly Biodiesel Production Survey"

U.S. Energy Information Administration | Monthly Biodiesel Production Report

Table 3. U.S. Inputs to Biodiesel Production (million pounds)

	Feedstock Inputs Vegetable Oils Animal Fats										
Period		Animal Fats									
erodīt.	Canola Oil	Corn Oil	Cottonseed Oil	Palm Oil	Soybean Oil	Other	Poultry	Tallow			
2011											
January	8	17	-	W	150	W	14	11			
February	26	13	•	W	150	W	14	11			
March	68	14	-	W	190	W	19	27			
April	88	20		w	236	W	15	47			
May	113	21	-	W	264	W	16	36			
June	75	34	-	W	311	W	23	49			
July	77	35	-	w	367	W	26	64			
August	84	37	W	w	398	W	34	38			
September	84	27	W	W	430	W	20	49			
October	69	30	W	w	527	w	31	39			
November	84	27	W	W	538	W	13	31			
December	71	30	W	W	592	W	15	27			
Total	847	304	w	w	4,153	w	240	431			
					4,133			431			
2012											
January	73	50	-	W	306	W	20	26			
February	90	51	-	W	321	W	8	20			
March	117	54	*	W	388	W	11	24			
April	112	54	5-14-1-1	W	350	W	13	45			
May	85	56		W	429	W	16	50			
June	103	55		W	369	w	14	28			
July	66	54		W	347	W	20	30			
August	48	59	_	W	385	W	21	44			
September	46	55	-	W	322	w	19	49			
October	16	59		W	307	w	14	44			
November	18	48		w	246	w	11	22			
December	16	50		W	273	w	9	3			
12 Month Total	790	646	•	W	4,042	w	176	385			
2013					***************************************						
lanuary	16	60		w	300	W	7	15			
ebruary	38	61		W	275	w	8	28			
March	39	71		w	424	W	9	53			
April	47	71		W	423	W	15	56			
May	w	91		W	416	W	20	61			
une	w	98		W	461	w	19	54			
uly	w	108		93	480	2	17	45			
August	79					the second second second second second					
September	W	106 95		W	510	2	17	48			
October	82	95 85		93	502		14	50			
LO Month Total	488			100	551	W 21	13	23			
to Month Total	400	846		476	4,342	31	139	433			
2012 10 Month Total	756	547		w	3 524	W	156	360			
2011 10 Month Total	692	248		W	3,524 3,023	the second second second second second	156 212	360 371			
= No data reported	032	240		VV	3,023	W	717	3/1			

^{- =} No data reported.

R = Revised

W = Withheld to avoid disclosure of individual company data.

⁽s) = Value is less than 0.5 of the table metric, but value is included in any associated total.

Totals may not equal the sum of components due to independent rounding.

Source: U.S. Energy Information Administration, Form EIA-22M "Monthly Biodiesel Production Survey"

U.S. Energy Information Administration | Monthly Biodiesel Production Report

Table 3a. Inputs to Biodiesel Production (continuation of table 3) (million pounds)

		Other Inputs						
Period	Anim	al Fats						
	White Grease	Other	Yellow Grease	Other	Algae	Other	Alcohol	Catalyst
2011								
January	31	3	18	10	-	7	20	4
February	48	2	19	11	-	4	20	3
March	60	12	33	10		7	34	5
April	55	8	52	11	-	(s)	46	7
May	57	10	41	15	*	(s)	48	8
June	55	7	50	18	•	1	58	9
July	49	6	45	20		2	62	9
August	48	15	47	19	-	5	63	9
September	40	4	50	21		(s)	64	9
October	29	7	42	21		(s)	71	10
November	35	8	39	20		(s)	71	10
December	26	5	35	20		(s)	68	11
12 Month Total	533	85	471	195		27	626	94
				193	· · · · · · · · · · · · · · · · · · ·	27	020	94
2012								
January	29	3	37	18	-	(s)	45	8
February	29	4	41	22		-	47	8
March	31	6	61	21		(s)	53	9
April	41	5	67	24			56	10
May	38	5	61	27		-	62	10
June	39	4	64	27		-	57	10
July	31	4	65	25	-	(s)	57	9
August	34	4	70	28		(s)	55	10
September	33	4	65	25	-		50	9
October	41	4	60	29		(s)	48	9
November	28	4	36	23	•		39	7
December	35	2	45	21		(s)	38	6
12 Month Total	408	48	670	289	•	1	607	105
2013								
January	31	3	46	23	-	1	50	7
February	32	3	51	23		(s)	47	7
March	36	2	72	15	-	(s)	63	10
April	48	2	79	19		3	70	11
May	41	W	88	20	-	W	78	12
lune	41	4	93	26	-	w	77	12
uly	43	2	97	31	-	W	90	13
August	41	W	92	31	_	W	89	13
September	38	W	107	22		w	89	13
October	36	1	81	27	-	W	81	13
10 Month Total	387	23	806	237	Zerosen-Kareka	38	734	111
1012 10 14								
2012 10 Month Total	346	43	591	246		1	530	92
= No data reported.	472	74	397	156		27	486	73

^{- =} No data reported.

R = Revised

W = Withheld to avoid discloaure of individual company data

⁽s) = Value is less than 0.5 of the table metric, but value is included in any associated total.

Totals may not equal the sum of components due to independent rounding.

Source: U.S. Energy Information Administration, Form EIA-22M "Monthly Biodiesel Production Survey"

U.S. Energy Information Administration | Monthly Biodiesel Production Report

Table 4. Biodiesel Producers and Production Capacity by State, October 2013

State	Number of Producers	Annual Production Capacity (million gallons per year)
Alabama	3	4
Alaska		
Arizona	1	
Arkansas	3	8
California	9	6.
Colorado		
Connecticut	3	. 1
Delaware		
District of Columbia		
Florida		
Georgia	3	
Hawaii		10
Idaho		
Illinois	1	
Indiana	5	16
	2	104
lowa	9	280
Kansas	1	
Kentucky	5	68
Louisiana	1	12
Maine	1	
Maryland		•
Massachussetts	1	
Michigan	2	18
Minnesota	4	10
Mississippi	3	109
Missouri	8	188
Montana	_	
Nebraska		
Nevada	1	i
New Hampshire	1	4
New Jersey		· · · · · · · · · · · · · · · · · · ·
New Mexico		
New York	1	C
North Carolina	4	10
North Dakota	1	85
Ohio	3	67
Oklahoma	1	15
Oregon	2	***************************************
Pennsylvania	4	18
Rhode Island		87
outh Carolina	12	1
outh Dakota	<u> </u>	40
ennessee	2	2
exas Itah	13	428
	1	10
ermont		
/irginia	3	9
Vashington	3	104
Vest Virginia		
Visconsin	3	29
Vyoming		
J.S. Total	112	2,190

^{- =} No data reported.

Totals may not equal the sum of components due to independent rounding.

Number of producers is a count of plants with operable capacity during the report month.

Source: U.S. Energy Information Administration, Form EIA-22M "Monthly Biodiesel Production Survey"

U.S. Energy Information Administration | Monthly Biodiesel Production Report

Table 5. Biodiesel (B100) Production by Petroleum Administration for Defense District (PADD) (million gallons)

	PADD									
Period	East Coast (PADD 1)	Midwest (PADD 2)	Gulf Coast (PADD 3)	Rocky Mountain (PADD 4)	West Coast (PADD 5)	U.S.				
2011										
January	3	30	1	-	1	35				
February	3	32	4	7)	1	40				
March	3	47	6		2	60				
April	3	54	10		3	71				
May	4	58	11	•	4	77				
June	4	56	14	•	7	81				
July	5	65	17		5	92				
August	5	66	20		5	95				
September	6	65	20		6	96				
October	7	73	22		4	105				
November	6	71	23	•	4	105				
December	6	77	23	3.5	4	109				
Total	56	695	171		46	967				
2012										
January	5	50	15	(s)	3	74				
February	4	54	18	(s)	3	79				
March	6	67	18	(s)	4	95				
April	6	63	21	1	3	94				
May	7	71	20	1	4	102				
June	6	67	17	(s)	3	93				
luly	5	66	15	(s)	3	89				
August	6	66	16	(5)	2	91				
September	6	60	14	(s)	3	82				
October	5	54	14	(s)	2	75				
November	2	44	8	(s)	2	57				
December	5	48	4	1	1	59				
Total	62	711	180	4	35	991				
2013										
anuary	4	52	9	(s)	1	66				
ebruary	4	50	12	(s)	1	68				
March	5	66	22	(s)	6	98				
April	6	70	23	(s)	7	106				
May	5	75	21	(s)	10	111				
une	6	79	22	(s)	5	113				
uly	6	82	30	(s)	8	128				
August	6	86	29	(s)	6	128				
September	6	81	30	(s)	9	127				
October	7	88	27	1	10	132				
10 Month Total	55	729	225	2	63	1,076				
2012 10 Month Total	56	618	168	3	30	874				
2011 10 Month Total	43	546	125		38	753				

^{- =} No data reported.

Totals may not equal the sum of components due to independent rounding.

B100 is the industry designation for pure biodiesel; a biodiesel blend contains both pure biodiesel and petroleum diesel fuel.

See Appendix A for a map of states included in each PADD.


Source: U.S. Energy Information Administration, Form EIA-22M "Monthly Biodiesel Production Survey"

U.S. Energy Information Administration | Monthly Biodiesel Production Report

R = Revised

⁽s) = Value is less than 0.5 of the table metric, but value is included in any associated total.

Appendix A Petroleum Administration for Defense Districts (PADD)



NOTE: MAP NOT TO SCALE

To:

Argyropoulos, Paul[Argyropoulos.Paul@epa.gov]

From:

Larry Schafer

Sent:

Tue 2/18/2014 5:37:53 PM FW: final version of presentation

Subject: FW: final version of presentation

2014.1.16 Draft Presentation on RFS Carbon Reduction - Final for Filing.pptx

For today's discussion ...

Larry Schafer

National Biodiesel Board

O: 202.737.8801

M: 202.997.8072

LSchafer@Biodiesel.org

Biodiesel - America's Advanced Biofuel!

www.americasadvancedbiofuel.com

1331 Pennsylvania Ave. NW

Suite 505

Washington DC 20004

From: David DeRamus [mailto:david.deramus@bateswhite.com]

Sent: Monday, January 27, 2014 9:34 AM

To: Larry Schafer

Cc: Anne Steckel; J. Alan Weber; Collin Cain

Subject: final version of presentation

Here's a final version of the PPT with the draft language taken off; the low 2014 forecast eliminated; a few

conforming word changes; and some minor format changes.

Thx

David

David DeRamus, Ph.D.

Managing Partner

direct: 202.216.1154 | mobile: 202.468.5604 | fax: 202.408.7838

1300 Eye Street NW, Suite 600, Washington, DC 20005

david.deramus@bateswhite.com

BATESWHITE.COM

Circulation of the Control of the Co

ATTENTION: This electronic message may contain PRIVILEGED and CONFIDENTIAL information and is intended for the use of the specific individual(s) and/or entity to which it is addressed. If you are not the intended recipient of this email, you are hereby notified that any unauthorized use, reliance, dissemination, or copying of this email, or the information contained in it or attached to it, is strictly PROHIBITED. If you have received this email in error, please delete it immediately and notify sender by reply email or by calling 202.408.6110.

******** ***** ATTACHMENT NOT DELIVERED **********

This Email message contained an attachment named image001.jpg

which may be a computer program. This attached computer program could contain a computer virus which could cause harm to EPA's computers, network, and data. The attachment has been deleted.

This was done to limit the distribution of computer viruses introduced into the EPA network. EPA is deleting all computer program attachments sent from the Internet into the agency via Email.

If the message sender is known and the attachment was legitimate, you should contact the sender and request that they rename the file name extension and resend the Email with the renamed attachment. After receiving the revised Email, containing the renamed attachment, you can rename the file extension to its correct name.

For further information, please contact the EPA Call Center at (866) 411-4EPA (4372). The TDD number is (866) 489-4900.

******* ***** ATTACHMENT NOT DELIVERED ****************

Standard **Biodiesel Renewable Fuel**

Analysis of Cost of Carbon Reduction DRAFT - PRELIMINARY ANALYSIS January 27, 2014

Summary of conclusions

- Biodiesel reduces CO2 emissions by 81% relative to diesel
 - Based on current mix of biodiesel feedstocks
 - Biodiesel carbon footprint steadily improving as mix of feedstocks changes
 - Biodiesel reduced 2013 CO2 emissions by 16.4 million metric tons
- Biodiesel production costs have declined, especially relative to petroleum diesel
 - Lower feedstock costs relative to petroleum diesel
 - Greater availability of feedstocks with lower carbon footprint
 - Technology changes allow greater use of lower-cost, lower-carbon feedstocks
- Cost of carbon reduction via biodiesel is currently \$71 \$91 per metric ton of CO2, before accounting for other benefits
- Biodiesel capacity and feedstocks are available to support further production increases with minimal cost/price impacts




Estimates of Biodiesel Cost of Carbon Reduction



Biodiesel has a significantly lower carbon footprint than petroleum

Biodiesel lifecycle carbon emissions, by feedstock



On average, biodiesel emits 81% less CO2 than petroleum diesel with current feedstocks



Currently, 103 gallons of biodiesel = 1 ton reduction in CO2 emissions


- With current feedstocks, biodiesel emits 81% less CO2 than petroleum diesel (normalized for energy content)
- 2013 wtd. average CO2 reduction: 21.3 pounds/gallon biodiesel
- 2204 lbs / 21.3 lbs/gallon = 103 gallons biodiesel to reduce CO2 emissions by 1 metric ton
- Current annual CO2 reductions due to biodiesel: 16.4 million metric tons
 - 1,700 million gallons x 21.3 lbs / 2204 = 16.43 million metric tons CO2e
 - Equivalent to the emissions from 3.4 million cars*

*Source: EIA's emissions calculator



Share of biodiesel production from lower carbon footprint feedstocks has increased, even with rising biofuel production

Biodiesel Use of Lowest Carbon Feedstock




Source: EIA Monthly Biodiesel Production Report (November 2013).

At the margin, increased biodiesel production has used distillers corn oil or waste grease as feedstocks



Changing feedstocks have made biodiesel increasingly effective at reducing carbon emissions


Gallons of Biodiesel Required to Reduce a Metric Ton of Carbon



Source: EIA Monthly Biodiesel Production Report , USEPA , USDA and CARB carbon reduction figures




Biodiesel production costs have been declining, especially relative to the price of petroleum diesel



Sources: Production cost estimates based on output and feedstock data from EIA for biodiesel and industry estimates for renewable diesel; feedstock prices from Jacobsen; processing costs from Iowa State biodiesel plant model.



The difference between the cost of biodiesel and the wholesale price of diesel has dropped by over 40% since 2011





Biodiesel production costs are falling with increased reliance on lower cost feedstocks


	2011	2012	2013
Weighted average feedstock cost	\$3.90	\$3.66	\$3.20
Average processing cost	\$0.67	\$0.66	\$0.69
Total production cost	\$4.57	\$4.32	\$3.89
Average wholesale diesel price	\$3.05	\$3.11	\$3.01
Biodiesel production cost difference	\$1.52	\$1.21	\$0.88

Sources: Production cost estimates based on output and feedstock data from EIA for biodiesel and industry estimates for renewable diesel; feedstock prices from Jacobsen; processing costs from lowa State biodiesel plant model.

Processing costs from lowa State model are likely overstated, especially with higher plant utilization in 2013



Lower biodiesel production costs reflect overall feedstock price decline and cost advantage of lower carbon footprint feedstocks



Source: Feedstock prices from Jacobsen; feedstock prices per pound were multiplied by 7.5 to determine the cost per gallon of biodiesel.




Expanded use of lower-cost/lower-carbon footprint feedstocks due to recent investments in new technology

- New technology increased production of lower carbon feedstocks
 - Distillers corn oil
 - Waste oils
- Updates to existing biofuel production systems
 - Flexible plants able to use different feedstock
- Alternative processing systems/pathways, e.g.:
 - Hydrotreating
 - Removes sulfur, oxygen, and nitrogen
 - Produces propane as a byproduct
 - Biomass-to-Liquid
 - Biomass is gasified
 - Resulting gas is converted into oil

Investments in technology for new feedstocks and more efficient biodiesel production due to increased biodiesel demand



Based on 2013 feedstocks and production costs, biodiesel's cost of carbon reduction is approximately \$71 - \$91 per metric ton of CO2



Changes in production costs and feedstocks have lowered biodiesel's cost of carbon reduction by 45% since 2011




Calculation details: CO2 reduction cost from biodiesel production cost differential relative to wholesale diesel price

- \$0.88 biodiesel cost differential relative to wholesale diesel
 - Wholesale diesel price in 2013 averaged \$3.01/gal
 - Production cost of biodiesel in 2013 was \$3.89/gal
- 21.3 lbs CO2 emission reduction from biodiesel displacing one gallon of petroleum diesel (0.00967MT/gal)
 - Based on carbon intensity estimates by feedstock and current feedstock production shares
- \$0.88/gal ÷ 0.00967MT/gal = \$91/MT CO2 reduction



An upper bound estimate of the cost of RFS2 is the cost differential between biodiesel and petroleum diesel



Blend Market with Volume Mandate

Biodiesel Market with Volume Mandate

Adapted from Taheripour and Tyner, "Welfare Assessment of the Renewable Fuel Standard: Economic Efficiency, Rebound Effect, and Policy Interaction in a General Equilibrium Framework," Purdue University Working Paper, June 2012, presented at 15th Annual Conference on Global Economic Analysis.



Cost of carbon reduction from biodiesel should be calculated net of other benefits

- RFS2 has three distinct policy objectives:
 - Agricultural policy: increase rural employment, farm incomes
 - "Energy independence:" reduce petroleum imports
 - Reduce GHG emissions
- Biodiesel is effective at reducing CO2 and other emissions, e.g. particulate matter (PM_{2.5})
 - Biodiesel reduces PM_{2.5} emissions by 47%¹
 - EPA estimate of benefits from PM_{2.5} reduction: \$230k 880k/ton²
- RFS2's other objectives also have significant monetary benefits


² EPA, "Report to Congress on Black Carbon," March 2012, p. 143.

Our initial estimate of biodiesel's cost of carbon reduction does not account for these other benefits



¹ EPA, "A Comprehensive Analysis of Biodiesel Impacts on Exhaust Emissions," October 2002, p. 37.

No indication that D4 RIN prices have had a significant impact on retail price of diesel fuel




Whether RIN costs are passed through to consumers is important in analyzing welfare effects of RFS2



Further increased biodiesel production is not constrained by available feedstocks


Biodiesel feedstock production forecast



Source: LMC International, "Current and Future Supply of RFS2 Qualifying and Non-qualifying Oils and Fats for Biofuels," July 2013.



Recent rapid growth in biodiesel has not increased soybean oil prices – additional production is unlikely to cause large feedstock cost increases




Source: Soy Oil prices from Jacobsen; RIN volumes from EPA EMTS database.



Increasing availability and recent price trends suggest incremental biodiesel production will come from lower carbon feedstocks

Biodiesel Use of Distillers Corn Oil as Feedstock




Sources: EIA Monthly Biodiesel Production Report, November 2013; LMC International Report, "Current and Future Supply of RFS2 Qualifying and Non-Qualifying Oils and Fats for Biofuels," July 2013.



Further increased biodiesel production is not constrained by available biodiesel capacity


2012 Biodiesel Production Capacity



Source: Biodiesel plant list 2-6-13 from Docket EPA-HQ-OAR-2013-0479; numbers are from a combination of NBB, EIA, and EPA databases.



D4 RIN prices are currently well below historical levels, reflecting growth in biodiesel supplies and production capacity



Source: Bloomberg; average of available indices.



Simulation models suggest biodiesel production can remain at 2013 levels or further expand without significantly increasing costs

	2013	2014 Forecast
Production, billion gallons	1.74	1.70
Weighted feedstock cost, \$/gal	\$3.20	\$2.70
Avg processing cost, \$/gal	\$0.50	\$0.50
Total Production Cost, \$/gal	\$3.70	\$3.20
Wholesale Diesel Price, \$/gal	\$3.01	\$2.92
Biodiesel cost differential, \$/gal	\$0.69	\$0.28
Cost of 1 metric ton CO2 reduction	\$71	\$29


Sources: 2013 production cost analysis with industry processing cost estimate; 2014 production cases from World Agricultural Economic and Environmental Services (WAEES) model.



Current biodiesel cost differential would be largely eliminated by a \$43/ton carbon price

- 2014 projected wholesale price of diesel = \$2.92/gallon
- A gallon of diesel fuel produces around 96g CO2e/MJ
- A gallon of diesel fuel has an energy intensity of 36.17 MJ/liter
- With carbon price of \$43/ton, diesel price would increase \$0.57/gal


$$2.92 \; \frac{\$}{gallon} + \; 96 \; \frac{g \; CO_2}{MJ} * \; 36.17 \; \frac{MJ}{liter} * \; 3.785 \\ \frac{liters}{gallon} * \; .000043 \; \frac{\$}{g \; CO_2} = \; 3.49 \; \frac{\$}{gallon}$$





Biodiesel cost of carbon reduction compares favorably to other policies

Cost of Reducing Carbon in Transportation Sector



Source for data regarding electric vehicle tax credit, hybrid vehicle tax credit, and CARS program: Ted Gayer, "Cash for Clunkers: An Evaluation of the Car Allowance Rebate System," *Economic Studies at Brookings*, October 2013.



3/9/2010 3/11/2010 3/11/2010 3/15/2010 3/17/2010 3/18/2010 3/25/2010 3/25/2010 3/30/2010 4/1/2010 4/6/2010 4/6/2010 4/8/2010 4/16/2010 4/16/2010 4/16/2010 4/16/2010 4/20/2010 4/20/2010 4/20/2010 4/20/2010 4/20/2010 4/20/2010 5/5/2010 5/6/2010 5/5/2010 5/11/2010 5/25/2010 5/26/2010 6/2/2010 6/2/2010 6/2/2010 6/2/2010 6/10/2010	0.175 0.175 0.175 0.185 0.19 0.205 0.225 0.255 0.245 0.277 0.266 0.275 0.255 0.245 0.245 0.255 0.245 0.255 0.335 0.315 0.35 0.31 0.36 0.31 0.36 0.35 0.31 0.36 0.35 0.31 0.36 0.35 0.395 0.41 0.41 0.4 0.36 0.35 0.395 0.41 0.41 0.4 0.36 0.35 0.395 0.41 0.41 0.4 0.36 0.35 0.395 0.41 0.41 0.4 0.36 0.35 0.395 0.41 0.41 0.4 0.36 0.35 0.395 0.305 0.305
6/2/2010 6/4/2010 6/8/2010	0.285 0.295 0.3075

6/18/2010 6/22/2010 6/23/2010 6/25/2010 6/25/2010 6/29/2010 7/1/2010 7/2/2010 7/8/2010 7/9/2010 7/12/2010 7/14/2010 7/14/2010 7/19/2010 7/21/2010 7/23/2010 7/23/2010 7/27/2010 7/28/2010 7/29/2010 7/30/2010 8/3/2010 8/5/2010 8/6/2010	0.325 0.3 0.35 0.335 0.485 0.51 0.505 0.4825 0.485 0.485 0.485 0.51 0.55 0.515 0.54 0.56 0.55 0.57 0.57 0.575 0.55 0.495 0.495 0.495 0.515
7/30/2010	0.535
8/3/2010	0.495
8/4/2010	0.495
9/17/2010	0.54
9/21/2010	0.57315
9/22/2010	0.57
9/24/2010	0.535
9/28/2010	0.54
9/30/2010	0.59
10/1/2010	0.6
10/5/2010	0.59

10/6/2010	0.56
10/7/2010	0.56
10/8/2010	0.55
10/12/2010	0.575
10/13/2010	0.575
10/18/2010	0.575
10/21/2010	0.56
10/22/2010	0.57
11/10/2010	0.61
11/11/2010 11/16/2010	0.6
11/17/2010	0.5975 0.605
11/18/2010	0.6112
11/19/2010	0.6137
11/23/2010	0.67
11/29/2010	0.69
12/1/2010	0.73
12/2/2010	0.775
12/3/2010	0.825
12/7/2010	0.83
12/8/2010 12/9/2010	0.93 0.825
12/3/2010	0.623
12/15/2010	0.5825
12/16/2010	0.56
12/17/2010	0.535
12/21/2010	0.535
12/22/2010	0.575
12/23/2010 12/24/2010	0.7175 0.7175
12/24/2010	0.7175
12/29/2010	0.735
12/31/2010	0.755
1/4/2011	0.79995
1/5/2011	0.77
1/6/2011	0.75
1/7/2011 1/12/2011	0.7675 0.7775
1/21/2011	0.7773
1/25/2011	0.88
1/26/2011	0.8775
1/28/2011	0.88
1/31/2011	0.9
2/1/2011	0.9225
2/4/2011 2/10/2011	0.935 1.05
2/14/2011	1.05
2/15/2011	1.21
2/16/2011	1.21
2/17/2011	1.235
2/22/2011	1.14
2/23/2011	1.005
2/25/2011	1.125

2/28/2011	1.1
3/1/2011	1.0625
3/2/2011	1.07
3/3/2011	1.0775
3/4/2011	1.08
3/7/2011	1.1075
3/8/2011	
	1.1125
3/9/2011	1.1125
3/10/2011	1.13
3/11/2011	1.1475
3/14/2011	1.145
3/15/2011	
	1.15
3/16/2011	1.0425
3/17/2011	1.0625
3/18/2011	1.13
3/21/2011	1.11
3/22/2011	1.085
3/23/2011	1.095
3/24/2011	1.115
3/25/2011	1.115
3/28/2011	1.12
3/29/2011	1.125
3/30/2011	1.18
3/31/2011	1.1825
4/1/2011	1.21
4/5/2011	1.255
4/6/2011	1.26
4/7/2011	1.255
4/8/2011	1.2775
4/11/2011	1.32625
4/12/2011	1.325
4/13/2011	1.345
4/14/2011	1.33
4/15/2011	1.32
4/18/2011	1.2925
4/19/2011	1.275
4/21/2011	1.2025
4/25/2011	1.275
4/26/2011	1.2625
4/27/2011	1.21
4/29/2011	1.18
5/2/2011	1.195
5/3/2011	1.25
5/4/2011	1.25
5/5/2011	1.255
5/6/2011	1.255
5/9/2011	1.27
5/10/2011	1.28
5/11/2011	1.24
5/12/2011	1.28
5/13/2011	1.27
5/16/2011	1.29
5/18/2011	1.32
	Vicination.

5/19/2011	1.33
5/20/2011	1.33
5/23/2011	1.34
5/24/2011	1.34
5/25/2011	1.295
5/26/2011	1.29
6/1/2011	1.32
6/2/2011	1.35
6/3/2011	1.35
6/6/2011	1.35
6/7/2011	1.37
6/8/2011	1.375
6/9/2011	1.41
6/13/2011	1.41
6/14/2011	1.405
6/15/2011	1.4
6/16/2011	1.375
6/17/2011	1.38
6/21/2011	1.345
6/22/2011	1.35
6/23/2011	1.37
6/24/2011	1.36
6/27/2011 6/28/2011	1.34 1.34
6/29/2011	1.34
6/30/2011	1.33
7/1/2011	1.34
7/5/2011	1.35
7/6/2011	1.335
7/7/2011	1.345
7/11/2011	1.345
7/12/2011	1.34
7/13/2011	1.34
7/14/2011	1.34
7/15/2011	1.33
7/18/2011	1.34
7/19/2011	1.33
7/20/2011	1.335
7/21/2011	1.325
7/22/2011	1.34
7/25/2011	1.3425
7/26/2011	1.36
7/27/2011	1.37
7/28/2011	1.37
7/29/2011	1.375
8/1/2011	1.37
8/2/2011	1.395
8/8/2011	1.4
8/9/2011 8/10/2011	1.405 1.45
8/11/2011	1.45
8/12/2011	1.49
8/15/2011	1.505
0, 10,2011	1.505

8/16/2011 8/17/2011 8/18/2011 8/19/2011 8/22/2011 8/23/2011 8/25/2011 8/25/2011 8/26/2011 8/29/2011 9/6/2011 9/6/2011 9/8/2011 9/9/2011 9/12/2011 9/14/2011 9/15/2011 9/16/2011 9/19/2011 9/20/2011 9/21/2011	1.54 1.635 1.695 1.695 1.685 1.685 1.685 1.685 1.685 1.685 1.685 1.895 1.895 1.897 1.97 1.975 1.92 1.89 1.875 1.75 1.75 1.75 1.75	
9/22/2011 9/23/2011	1.695 1.72	
9/26/2011	1.71	
9/27/2011	1.67	
9/28/2011 9/29/2011	1.605 1.61	
9/30/2011	1.55	
10/4/2011	1.495	
10/5/2011	1.535	
10/6/2011	1.535	
10/11/2011	1.475	
10/13/2011	1.44	
10/14/2011	1.44	
10/17/2011	1.38	
10/18/2011 10/19/2011	1.38 1.375	
10/20/2011	1.325	
10/21/2011	1.255	
10/24/2011	1.17	
10/25/2011	1.17	
10/26/2011	1.06	
10/27/2011 10/28/2011	1.06 1.225	
10/20/2011	1.16	
11/1/2011	1.16	
11/2/2011	1.14	
11/3/2011	1.14	
11/4/2011	1.14	
11/7/2011 11/8/2011	1.21 1.21	
11/9/2011	1.4	
. 11012011	1.7	

11/11/2011	1 15
	1.45
11/14/2011	1.395
11/15/2011	1.395
11/16/2011	1.5
11/17/2011	1.5
11/18/2011	1.5
11/21/2011	1.44
11/22/2011	1.44
11/23/2011	1.435
11/28/2011	1.42
11/29/2011	1.42
11/30/2011	1.375
12/1/2011	1.435
12/2/2011	1.38
12/5/2011	1.32
12/6/2011	1.32
12/7/2011	
	1.33
12/8/2011	1.305
12/9/2011	1.38
12/12/2011	1.3
12/13/2011	1.3
12/14/2011	1.295
12/15/2011	1.2
12/16/2011	1.2
12/19/2011	1.23
12/20/2011	1.255
12/21/2011	1.27
12/27/2011	1.275
12/28/2011	1.28
12/29/2011	1.25
12/30/2011	1.245
1/3/2012	1.3875
1/4/2012	1.39
1/5/2012	1.4
1/6/2012	1.4
1/9/2012	1.4525
1/10/2012	1.445
1/11/2012	1.4475
1/12/2012	1.4125
1/13/2012	1.43
1/17/2012	1.42
1/18/2012	1.425
1/20/2012	1.42
1/23/2012	1.4025
1/24/2012	1.4
1/25/2012	1.3975
1/26/2012	1.395
1/30/2012	1.365
1/31/2012	1.365
2/1/2012	1.3425
2/2/2012	1.355
2/3/2012	1.4175
2/13/2012	1.385

2/14/2012	1.415
2/15/2012	1.42
2/22/2012	1.435
2/23/2012	1.43
2/24/2012	1.425
2/27/2012	1.41
2/28/2012	1.435
2/29/2012	1.43
3/1/2012	1.415
3/2/2012	1.4
3/5/2012	1.39
3/6/2012	1.38375
3/7/2012	1.3675
3/8/2012	1.325
3/9/2012	1.3375
3/12/2012	1.3125
3/13/2012	1.33
3/14/2012	1.3375
3/15/2012	1.3325
3/16/2012	1.3325
3/19/2012	1.335
3/20/2012	1.33
3/21/2012	1.345
3/22/2012	1.3325
3/27/2012	1.3425
3/28/2012	1.3275
4/3/2012	1.3175
4/5/2012	1.305
4/9/2012	1.3325
4/10/2012	1.345
4/11/2012	1.3325
4/12/2012	1.3425
4/13/2012	1.3325
4/16/2012	1.315
4/17/2012	1.32
4/18/2012	1.335
4/19/2012	1.34
4/20/2012	1.335
4/23/2012	1.3225
4/24/2012	1.3375
4/25/2012	1.3275
4/26/2012	1.325
4/27/2012	1.3225
5/1/2012	1.34
5/2/2012	1.36
5/3/2012	1.36
5/4/2012	1.355
5/7/2012	1.38
5/8/2012	1.3725
5/9/2012	1.3725
5/14/2012	1.3625
5/16/2012	
	1.35
5/17/2012	1.31

5/18/2012	1.35
5/21/2012	1.34
5/22/2012	1.3425
5/24/2012	1.3423
5/30/2012	
	1.3175
5/31/2012	1.3125
6/7/2012	1.11
6/8/2012	1.155
6/11/2012	1.225
6/12/2012	1.22
6/13/2012	1.205
6/14/2012	1.21
6/15/2012	1.2
6/18/2012	1.19
6/19/2012	1.18
6/20/2012	1.17
6/21/2012	1.17
6/22/2012	1.145
6/25/2012	1.135
6/26/2012	1.14
6/27/2012	1.16
7/5/2012	1.2
7/6/2012	1.2
7/9/2012	1.21
7/10/2012	1.215
7/11/2012	1.2
7/12/2012 7/13/2012	1.24 1.245
7/16/2012	1.245
7/19/2012	1.2375
7/20/2012	1.2425
7/23/2012	1.23875
7/24/2012	1.225
7/27/2012	1.2025
7/30/2012	1.185
7/31/2012	1.165
7/27/2012	1.17
7/30/2012	1.17
7/31/2012	1.1
8/1/2012	1.1
8/2/2012	1.1
8/3/2012	1.11
8/6/2012	1.1
8/7/2012	1.1
8/13/2012	1.05
8/14/2012	1.05
8/15/2012	1.02
8/16/2012	1
8/20/2012	0.975
8/21/2012	0.965
8/22/2012	0.975
8/27/2012	0.99
8/28/2012	1.02

8/29/2012	1.005
8/30/2012	0.995
9/4/2012	1.02
9/5/2012	1
9/6/2012	1
9/7/2012	0.98
9/10/2012	0.98
9/11/2012	1
9/12/2012	0.995
9/14/2012	0.99
9/17/2012	0.985
9/19/2012	0.975
9/20/2012	0.94
9/24/2012	0.885
9/25/2012	0.8
9/26/2012	0.78
9/27/2012	0.765
10/1/2012	0.74
10/2/2012	0.69
10/5/2012	0.55
10/8/2012	0.505
10/10/2012	0.435
10/15/2012	0.45
10/16/2012	0.55
10/17/2012	0.69
10/19/2012	0.62
10/22/2012	0.6
10/24/2012	0.5
10/25/2012	0.5
10/29/2012	0.5
10/30/2012	0.51
11/1/2012	0.61
11/5/2012	0.55
11/6/2012	0.54
11/7/2012	0.52
11/8/2012	0.56167
11/9/2012	0.57333
11/12/2012	0.56667
11/13/2012	
	0.48
11/16/2012	0.505
11/19/2012	0.5
11/26/2012	0.57167
11/27/2012	0.57
11/28/2012	0.55833
11/29/2012	0.54333
11/30/2012	0.54333
12/3/2012	0.55
12/4/2012	0.56333
12/5/2012	0.55833
12/6/2012	0.54667
12/10/2012	0.56167
12/11/2012	0.56
12/12/2012	0.59

12/13/2012	0.58667
12/28/2012	0.62667
12/31/2012	0.62667
1/2/2013	0.395
1/3/2013	0.38
1/7/2013	0.3725
1/8/2013	0.3725
1/9/2013	0.475
1/10/2013	0.475
1/14/2013	0.445
1/15/2013	0.49
1/16/2013	0.49
1/17/2013	0.49
1/18/2013 1/21/2013	0.49
1/21/2013	0.49 0.49
1/23/2013	0.49
1/24/2013	0.49
1/25/2013	0.47
1/28/2013	0.4625
1/29/2013	0.4675
1/30/2013	0.495
1/31/2013	0.495
2/1/2013	0.495
2/11/2013	0.5375
2/12/2013	0.4925
2/13/2013	0.4925
2/14/2013	0.4925
2/15/2013 2/19/2013	0.4925
2/21/2013	0.515 0.595
2/22/2013	0.5775
2/25/2013	0.545
2/27/2013	0.64
2/28/2013	0.63
3/1/2013	0.67
3/4/2013	0.73
3/5/2013	0.73
3/6/2013	0.765
3/7/2013	0.825
3/8/2013	0.995
3/11/2013	0.9825
3/12/2013 3/13/2013	0.925
3/14/2013	0.765
3/15/2013	0.75
3/18/2013	0.76
3/19/2013	0.76
3/20/2013	0.735
3/21/2013	0.735
3/25/2013	0.7025
3/26/2013	0.745
4/1/2013	0.7225

4/2/2013 4/3/2013 4/4/2013 4/8/2013 4/9/2013 4/10/2013 4/12/2013 4/12/2013 4/19/2013 4/23/2013 4/25/2013 4/25/2013 4/26/2013 5/2/2013 5/2/2013 5/3/2013 5/6/2013 5/8/2013 5/1/2013 5/1/2013 5/1/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/15/2013 5/20/2013 6/3/2013 6/5/2013 6/5/2013 6/12/2013	0.72 0.735 0.745 0.89 0.92 0.855 0.8275 0.79 0.83 0.85 0.83 0.82 0.81 0.82 0.81 0.82 0.83 0.94 0.9 0.845 0.855 0.87 0.855 0.87 0.855 0.87 0.855 0.89 0.915 0.935 0.945 0.945 0.945 0.945 0.945 0.955 0.955 0.965 0.995
6/12/2013 6/13/2013 6/14/2013	0.95 0.965
6/21/2013	0.995

6/24/2013 6/25/2013 6/26/2013 6/27/2013 6/28/2013 7/1/2013 7/3/2013 7/8/2013 7/9/2013 7/10/2013 7/11/2013 7/15/2013 7/17/2013 7/17/2013 7/18/2013 7/19/2013 7/19/2013 7/22/2013 7/22/2013 7/24/2013 7/25/2013	0.97 0.985 1.005 1.0125 1.045 1.03 1.025 1.055 1.0825 1.165 1.145 1.195 1.2875 1.35 1.44 1.415 1.355 1.3475 1.27
7/2/2013	1.03
7/3/2013	1.025
7/8/2013	1.055
7/10/2013	1.165
7/11/2013	1.145
7/12/2013	1.195
7/17/2013	1.44
7/18/2013	1.415
7/19/2013	1.415
7/23/2013	1.3475
7/24/2013	1.27
7/25/2013	1.005
7/26/2013	1.005
7/29/2013	0.975
7/30/2013	1.075
7/31/2013	1.155
8/1/2013	1.115
8/5/2013	1.095
8/6/2013	0.975
8/7/2013	0.88333
8/8/2013	0.78
8/9/2013	0.76
8/12/2013	0.82
8/13/2013	0.83
8/15/2013	0.91
8/16/2013	0.91667
8/19/2013	0.92667
8/20/2013	0.87667
8/21/2013	0.86667
8/22/2013	0.83
8/23/2013	0.82667
8/26/2013	0.83667
8/27/2013	0.81667
8/28/2013	0.78667
8/29/2013	0.76667
8/30/2013	0.76667
9/3/2013	0.84
9/4/2013	0.81667
9/5/2013	0.80667
9/6/2013	0.78667
9/9/2013	0.79667
9/10/2013	0.78667
9/11/2013	0.74667

9/12/2013	0.73667
9/13/2013	0.72333
9/16/2013	0.69333
9/17/2013	0.61333
9/18/2013	0.64333
9/19/2013	0.70333
9/20/2013	0.69333
9/23/2013	0.66333
9/24/2013	0.66333
9/25/2013	0.67333
9/26/2013	0.68333
9/27/2013	0.63333
9/30/2013	0.60333
10/1/2013	0.63333
10/2/2013	0.58333
10/3/2013	0.62333
10/4/2013	0.64333
10/7/2013	0.63333
10/8/2013	0.62333
10/9/2013	0.65333
10/10/2013	0.65333
10/11/2013	0.54333
10/14/2013	0.54
10/15/2013	0.57333
10/16/2013	0.55333
10/17/2013	0.52667
10/18/2013	0.52667
Delighan Telephone probability of the control of th	
10/21/2013	0.52333
10/22/2013	0.51333
10/23/2013	0.46333
10/24/2013	0.41333
10/28/2013	0.30333
10/30/2013	
	0.35667
11/6/2013	0.42333
11/7/2013	0.41333
11/8/2013	0.38333
11/11/2013	0.36333
11/12/2013	
	0.31
11/13/2013	0.29333
11/14/2013	0.35333
11/15/2013	0.28667
11/18/2013	0.27667
11/19/2013	0.26667
11/20/2013	0.315
11/21/2013	0.30333
11/22/2013	0.32333
11/26/2013	0.33
11/27/2013	0.33
To	

Sales

Registered fa 2.4
Registered fa 0.4
Facilities that 0.8

Year	ConsumpticP	rojected Production	Distillers	Cmetric tonnes
2011	138	377	304	137.93103
2012	293	538	646	293.10345
2013	500	961	1102	500
2	014	1419		
2	015	1536		

Best Ava	ailable	
regular diese	27.9	
canola	12.9	
soy	6.2	
used cooking	3.7	
other recycle	3.7	
animal fat	3.7	
distillers corr	1.2	

	USDA/Idaho	0.764 soy	89.7	96	27.91197
soy	0.5 USDA/Idaho	Osoy	96	96	27.91197
corn oil	0.1 CARB 2011	0.95833 corn oil	96	4	1.16300
used cooking oil	0.1 USEPA 201(0.86919 waste grea	96	12.558	3.65123
other recycled grease	0.03 USEPA 201(0.86919 waste grea	96	12.558	3.65123
animal fat	0.13 USEPA 201(0.86919 waste grea	96	12.558	3.65123
renewable diesel	0.1 CARB 2011	0.79531 tallow RD	96	19.65	5.71323
canola	0.04 USEPA 2010	0.53748 canola	96	44.4015	12.90972

Series 1	
Electric vehicle tax credit	300
Hybrid vehicle tax credit	177
CARS program	91
Biodiesel	71

Year	Gallons / Metric ton of Carbon	GHG reduction in lbs CO2e/gallon of biodiesel	Conversion to Tons	Gallons / Ton
2011	109	20	0.0091	6109.13973
2012	107	20.	7 0.0093	9106.50351
2013	104	21	0.0096	6103.50341

Year	Biodiesel ProductiorD	istillers	CoBiodiesel Production2	Soybean a Co	orn, Tallov	v, Yellow Grease a	а
2011	1,122,699,193.00	34%	,122,699,193	5000	2549	0.33769	
2012	1,146,899,177.00	40%	1146899177	4832	3218	0.39978	
2013	1,740,915,464.00	47%	1740915464	4131	3722	0.47398	

nd other Oils

2663

year	2011	2012	2013
Production Cost of Biodiese	\$ 4.5	7 \$ 4.32	\$ 3.89
Wholesale Price of Diesel	\$ 3.0	5 \$ 3.11	\$ 3.01
Difference	\$ 1.5	2 \$ 1.21	\$ 0.88

Based on Iowa StaBased on NBB processing cost estimates

2011	\$ 166	\$ 158
2012	\$ 128	\$ 121
2013	\$ 91	\$ 71

Soybean Oi Distiller's CcColumn1

 2011
 4.05598
 2.98403

 2012
 3.87813
 2.92350

 2013
 3.65178
 2.65779

To resize chart data range, drag lower right corner of range.

2666

year	2011	2012	20	13
Production Cost of Biodiesel	\$ 4.5	7 \$	4.32	\$ 3.89
Wholesale Price of Diesel	\$ 3.0	5 \$	3.11	\$ 3.01
Based on Iowa State processing cost estimate	\$ 1.5	2 \$	1.21	\$ 0.88
Based on NBB processing cost estimates	\$ 1.4	5 \$	1.15	\$ 0.69